

Two complete finitary sequent calculi for reflexive common knowledge

Aurimas Paulius Girčys, Regimantas Pliuškevičius

Vilnius University, Institute of Mathematics and Informatics

Akademijos 4, LT-08663 Vilnius

E-mail: aurimas.gircys@gmail.com, regimantas.pliuskevicius@mii.vu.lt

Abstract. This paper discusses the use of complete sequent calculi for reflexive common knowledge logic. Description of language and complete infinitary calculus for RCL is presented. Then finitary calculi RCL_I and RCL_L are introduced and completeness of finitary calculi RCL_I and RCL_L is proven.

Keywords: common knowledge logic, reflexive common knowledge logic, sequent calculi.

1 Introduction

A reflexive common knowledge logic (RCL) containing individual knowledge operators, reflexive “common knowledge” and “everyone knows” operators is considered. Complete sequent calculi for reflexive common knowledge logic is discussed, finitary calculi RCL_I and RCL_L are introduced and completeness of these calculi is obtained using completeness of the infinitary calculus for RCL.

2 Description of language and complete infinitary calculus for RCL

The language of considered RCL contains a set of propositional symbols $P, P_1, P_2, \dots, Q, Q_1, Q_2, \dots$ the set of logical connectives $\supset, \wedge, \vee, \neg$; finite set of agent constants i, i_1, i_2, \dots ; multiple knowledge modality $K(i)$, where i is an agent constant; everyone knows operator E ; common knowledge operator C .

A formula of RCL is defined inductively as follows: every propositional symbol is a formula; if A, B are formulas, then $(A \supset B)$, $(A \wedge B)$, $(A \vee B)$, $\neg(A)$ are formulas; if i is an agent, A is a formula, then $K(i)A$ is a formula; if A is a formula, then $E(A)$ and $C(A)$ are formulas. The operator $K(i)$ behaves as modality of multi-modal logic K_n [1].

The formula $K(i)A$ means “agent i knows A ”. The formula $E(A)$ means “every agent knows A ”, i.e. $E(A) = \bigwedge_{i=1}^n K(i)A$ (n is a number of agents). The formula $C(A)$ means “ A is common knowledge of all agents”; it is assumed that there is perfect communication between agents. The operator C and E behave as modalities of modal logic $S5$. In addition these operators satisfy the following powerful properties: $C(A) = A \wedge E(C(A))$ (fixed point) and $A \wedge C(A \supset E(A)) \supset C(A)$ (induction). Formal semantics of the formulas $K(i)$, $E(A)$, $C(A)$ are defined as in the reflexive common knowledge logic [3].

Bellow we consider calculi based on sequents, i.e., formal expressions $A_1, \dots, A_k \rightarrow B_1, \dots, B_m$, where $A_1, \dots, A_k (B_1, \dots, B_m)$ is a multiset of arbitrary formulas. The infinitary calculus, denoted by RCL_ω , for RCL is defined by following postulates [3].

Axiom: $\Gamma, A \rightarrow \Delta, A$.

Rules consist of logical rules and modal ones. Logical rules consist of traditional invertible rules for logical symbols.

Modal rules:

$$\frac{\Gamma \rightarrow A}{\Pi, K_i \Gamma \rightarrow \Delta, K_i(A)} (K_i),$$

where $K_i \Gamma = K_i A_1, \dots, K_i A_n$ ($n \geq 0$); Π, Δ consist of multisets of arbitrary formulas.

$$\frac{\bigwedge_{i=1}^m K_i(A), \Gamma \rightarrow \Delta}{E(A), \Gamma \rightarrow \Delta} (E \rightarrow),$$

$$\frac{\Gamma \rightarrow \Delta, \bigwedge_{i=1}^m K_i(A)}{\Gamma \rightarrow \Delta, E(A)} (\rightarrow E),$$

where m is number of agents.

$$\frac{A, E(C(A)), \Gamma \rightarrow \Delta}{C(A), \Gamma \rightarrow \Delta} (C \rightarrow),$$

$$\frac{\Gamma \rightarrow \Delta, A; \Gamma \rightarrow \Delta, E(A); \dots; \Gamma \rightarrow \Delta, E^k(A) \dots}{\Gamma \rightarrow \Delta, C(A)} (\rightarrow C_\omega),$$

where $E^0(A) = A; E^k(A) = E(E^{k-1}(A)), k \geq 1$.

It is known (see e.g. [3]) that calculus RCL_ω is sound and complete.

$$\frac{\Gamma \rightarrow A}{\Pi, E(\Gamma \rightarrow \Delta, E(A))} (E).$$

The rule is derivable in RCL^* where RCL^* is obtained from RCL_ω by dropping the rule $(\rightarrow C_\omega)$.

Let $\Gamma = A_1, \dots, A_n$ then derivability of (E) is carried out in the following way:

$$\frac{\frac{\frac{A_1, \dots, A_n \rightarrow A}{\dots, K_i(A_1), \dots, K_i(A_n) \dots \rightarrow \dots K_i(A) \dots} (K_i)}{\Pi, E(A_i), \dots, \bigwedge_{i=1}^m K_i(A_i), \dots, E(A_n) \rightarrow \bigwedge_{i=1}^m K_i(A)} (\wedge \rightarrow)(\rightarrow \vee)}{\Pi, E(A_1), \dots, E(A_n) \rightarrow E(A)} (E \rightarrow), (\rightarrow E).$$

Derivations in RCL_ω are built in the form of the infinite tree, each branch of this tree is finitary. The height of a derivation D (denoted by $O(D)$) is evaluated in ordinals.

A derivation D in RCL_ω is called atomic if all axioms occurring in D are the form $\Gamma, P \rightarrow \Delta, P$.

Lemma 1. *An arbitrary derivation in RCL_ω may be transformed into an atomic one.*

Proof. Let us denote by $g(A)$ the complexity of A defined by the number of occurrences of logical and knowledge operators C, E, K_i in A . The lemma is proved by induction on $g(A)$.

It is easy to see that all rules of RCL_ω , except (K_i) are invertible. Let us present a specialization of the rule (K_i) which is existential invertible.

A sequent S is a primary if $S = \Sigma_1, K(\Gamma_1) \rightarrow \Sigma_2, K(\Gamma_2)$, where Σ_i ($i \in \{1, 2\}$) is empty or consists of propositional symbols, $K(\Gamma_i)$ ($i \in \{1, 2\}$) is empty or consists of formulas of the shape $K_l(A)$.

Lemma 2. *By backward applications of rules, except (K_i) of RCL_ω any sequent S can be reduced to a set of primary sequents S_1, \dots, S_n ($n \geq 1$) such that if $RCL_\omega \vdash S_l$ then $\forall l$ ($l \geq 1$) $RCL_\omega \vdash S_l$.*

Proof. Follows from invertibility of rules RCL_ω except (K_i) .

Let RCL'_ω be the calculus obtained from RCL_ω replacing the rule (K_i) by the following one:

$$\frac{\Gamma_p \rightarrow A}{\Sigma_1, K_1^a(\Gamma_1), \dots, K_n^a(\Gamma_n) \rightarrow \Sigma_2, K_1^s(\Delta_1), \dots, K_l^s(A), \dots, K_m^s(\Delta_m)} (K'_i),$$

$n \geq 0, m \geq 0; K_l^a = K_p^s, \Sigma_1 \cap \Sigma_2 = \emptyset$.

Lemma 3 [Existential invertability of the rule (K'_i) in RCL_ω]. *Let $S = \Sigma_1, K_1^a(\Gamma_1), \dots, K_n^a(\Gamma_n) \rightarrow \Sigma_2, K_1^s(\Delta_1), \dots, K_m^s(\Delta_m)$ be a primary sequent satisfying the condition of the conclusion of (K'_i) and let $RCL_\omega \vdash^D S$, then there exists a formula $K_l^s(A)$ such that $RCL_\omega \vdash \Gamma_p \rightarrow A$.*

Proof. From Lemma 1 it follows that all axioms in D are atomic ones. Another hand, $\Sigma_1 \cap \Sigma_2 = \emptyset$ therefore $h(D) > 1$. Therefore from the scope of the rule (K'_i) it follows that there exists a formula $K_l^s(A)$ from the succedent of S such that $RCL_\omega \vdash \Gamma_p \rightarrow A$.

3 Finitary calculi RCL_I and RCL_L

Infinitary calculus RCL_ω possesses the following beautiful property: it allows to present simple and evident completeness proof (see e.g. [3]). Despite of this property: all derivation containing infinitary rule $(\rightarrow C_\omega)$ are informal. To avoid this bad property several finitary complete sequent calculi for RCL can be presented [2].

(1) Calculus containing invariant-like rule.

The finitary calculus RCL_I is obtained from the calculus RCL_ω replacing infinitary rule $(\rightarrow C_\omega)$ by following (cut) – like rule:

$$\frac{\Gamma \rightarrow \Delta, I; I \rightarrow E(I); I \rightarrow A}{\Gamma \rightarrow \Delta, C(A)} \rightarrow (C_I)$$

where the formula I (called an invariant formula) is constructed from subformulas of formulas in the conclusion of the rule. There are some works in which constructive methods for finding invariant formulas in sequent calculi of epistemic logic are presented, e.g. [4, 5]. Using these methods we can find invariant formulas and for the rule $(\rightarrow C_I)$.

(2) Calculus containing weak-induction like rule and loop axiom.

The finitary calculus RCL_L is obtained from the calculus RCL_I in the following way:

- (a) replacing the invariant rule $(\rightarrow C_I)$ by the following rule:

$$\frac{\Gamma \rightarrow \Delta, A; \Gamma \rightarrow \Delta, E(C(A))}{\Gamma \rightarrow \Delta, C(A)} (\rightarrow C_L).$$

This rule corresponds to the so-called weak-induction axiom: $A \wedge E(C(A)) \supset C(A)$.

- (b) Adding loop-type axioms as follows: a sequent S' is a loop type axiom if (1) S' is above a sequent S on a branch of derivation tree, (2) S' is such that it subsumes S' ($S \succcurlyeq'$ in notation), i.e. we can get S' from S using structural rules of weakening and contraction, in separate case $S = S'$.
- (c) There is right premise of $(\rightarrow C_L)$ between S and S' .

The completeness of finitary calculi RCL_I and RCL_L is obtained proving that the calculi RCL_ω , RCL_I and RCL_L are equivalent to each other. The completeness of RCL_ω is used.

References

- [1] P. Aleate and R. Gore. Cut-free single pass tableaux for the logic of common knowledge. In *Proc. of Workshop on Agents and Deduction at Tableaux 2007*, 2007.
- [2] K. Brunlez and T. Studez. Symantic cut-elimination for common knowledge. In *Proc. of Methods for Modalities, vol. 5*, 2007.
- [3] J.J.Ch. Meyer and W. von der Hoek. *Epistemic Logic for AI and Computer Science*. Cambridge University Press, Cambridge, 1995.
- [4] R. Pliuškevičius. The saturation tableaux for linear miniscoped horn-like temporal logic. *J. Autom. Reas.*, **13**:51–67, 1994.
- [5] M.K. Valiev. Decision complexity of variants of propositional dynamic logic. In P. Dembinski(Ed.), *Proc. MFCS'80*, LNCS, vol. 88, pp. 656–664. Springer-Verlag, Berlin, 1980.

REZIUMĖ

Du pilni finitariniai skaičiavimai refleksyviai bendrojo žinojimo logikai

P. Girčys, R. Pliuškevičius

Straipsnyje pateikiami du pilni sekvenciniai skaičiavimai bendrojo žinojimo logikai. Pristatyta kalba ir pilnas begalinis skaičiavimas skirtas RCL. Straipsnyje pristatyti baigtiniai skaičiavimai RCL_I ir RCL_L , ir įrodytas tų skaičiavimų pilnumas remiantis baigtinio skaičiavimo RCL_ω pilnumu.

Raktiniai žodžiai: bendrojo žinojimo logika, refleksyvi bendrojo žinojimo logika, sekvencinis skaičiavimas