

Sheaves on quantaloids

Remigijus Petras GYLYS (MII)

1. Introduction

This note considers sets valued in possibly non-unital quantaloids (categories enriched in the category of complete sup-lattices subject to certain laws). Quantaloids are a natural and useful categorical generalization of quantales (which are the one-object quantaloids, i.e., complete sup-lattices equipped with a multiplication). In the "logical" approach (see [3]) a sheaf on a topological space U corresponds to the "complete" set valued in the quantale $O(U)$ of open sets of U with the intersection as a multiplication. This approach to sheaves was further developed by U. Höhle [4,5] (based on a symmetric or "right symmetric" (but non-idempotent) quantale) and by F. Borceux and G. van den Bossche [1], C.J. Mulvey and M. Nawaz [6] (based on an idempotent quantale). In the present note, we provide a setting for sheaves on quantaloids (which is more general than ones mentioned above) taking our inspiration in G. van den Bossche's work [2] where sets valued in quantaloids are presented using "matrices". Our results are submitted without proofs. We are going to detail it in a subsequent paper.

2. Preliminaries on quantaloids and matrices over their

We begin by reviewing a few permanent definitions.

DEFINITION 2.1. A *quantaloid* is a locally small category Q (not necessarily having units) such that:

- (i) for all u, v objects in Q , the hom-set $Q(u, v)$ is a complete lattice,
- (ii) composition of morphisms of Q (in this note denoted by $\&$) preserves arbitrary joins in both variables:

$$p \& \bigvee_i q_i = \bigvee_i p \& q_i \text{ and } (\bigvee_i p_i) \& q = \bigvee_i p_i \& q$$

for all morphisms p, q of Q and for all families $(p_i), (q_i)$ of morphisms of Q (forming respective composable pairs).

Note that we use the unconventional left-to-right direction for composition of morphisms. Examples of the one-object quantaloids (which are called quantales) include frames (and thus

complete Boolean algebras) and various ideal lattices of rings or C^* -algebras. Many other examples of quantales and quantaloids can be found in [1]–[8].

From now Q will be an arbitrary quantaloid (not necessarily having units) having a small set of objects. Let Q_0 denote this set and Q_1 the set of morphisms of Q . Let $Sets/Q_0$ denotes the category whose objects are families X of sets X_u indexed by $u \in Q_0$. An element $x \in X_u$ will be called an element over u and we shall sometimes write $d(x)$ for u and $x \in X$ for $x \in X_u$. Morphisms in $Sets/Q_0$ are families of maps $f_u : X_u \rightarrow Y_u$.

DEFINITION 2.2 (Definition 1.3 [2]). Let Q be a quantaloid and X and Y be two objects of $Sets/Q_0$. A matrix M from X to Y assigns to each pair x, y of $X \times Y$ an element of Q_1 : $m_{x,y} : d(x) \rightarrow d(y)$. Matrices compose by "matrix multiplication": for $M : X \rightarrow Y$, and $N : Y \rightarrow Z$, the composite $M \& N = L : X \rightarrow Z$ has its general element given by

$$l_{x,z} = \bigvee_{y \in Y} m_{x,y} \& n_{y,z}.$$

3. Q -sets and bimodules

The notions of a Q -set and of a bimodule which will be given in this section are taken from [2].

DEFINITION 3.1 (Definition 2.1 [2]). Let Q be a quantaloid. A Q -set is an object X of $Sets/Q_0$ provided with a matrix $A : X \rightarrow X$ satisfying the following:

Idempotency: $A \& A = A$

A Q -set (X, A) will be called *separated* whenever it satisfies

Separation: if $a_{x,x''} = a_{x',x''}$ and $a_{x'',x} = a_{x'',x'}$ for all $x'' \in X$, then $x = x'$.

An element $x \in X$ of a Q -set (X, A) will be said to be *strict* provided that it satisfies

Strictness: $a_{x,x} \& a_{x,x'} = a_{x,x'}$ and $a_{x',x} \& a_{x,x} = a_{x',x}$ for all $x' \in X$ and a Q -set (X, A) itself will be called *strict* whenever every element $x \in X$ is strict.

We shall usually write a_x for $a_{x,x}$. Note that strict or separated Q -sets were not considered by G. van den Bossche in [2]. Conditions corresponding to Separation appear in [3, 4, 5]. Q -sets as defined in [3.6] are strict Q -sets in our sense.

DEFINITION 3.2 (Definition 2.6 [2]). Let (X, A) and (Y, B) be Q -sets. A *bimodule* (or morphism as called in [2]) \mathcal{F} from (X, A) to (Y, B) , written $\mathcal{F} : (X, A) \rightarrow (Y, B)$, is a pair of adjoint matrices $F : X \rightarrow Y$, $F^\# : Y \rightarrow X$, $F \dashv F^\#$, compatible with the structural matrices A and B , i.e., a pair of matrices satisfying the following:

Compatibility: $F = A \& F = F \& B$ and $F^\# = B \& F^\# = F^\# \& A$,

Adjunction: $A \leq F \& F^\#$ (unit) and $F^\# \& F \leq B$ (counit).

Bimodules compose just by composition of matrices. Structural matrices are their own adjoints and determine the units for bimodule composition. Thus Q -sets and their bimodules constitute a category denoted by Q -Sets. Our first result is the following

PROPOSITION 3.3 (cf. Corollary 13 [6]). Given a bimodule $\mathcal{F} = (F, F^\#) : (X, A) \rightarrow (Y, B)$ between Q -sets of a quantaloid Q ,

(i) if $A \geq F \& F^\#$, then \mathcal{F} is a monic in Q -Sets (i.e., F can be ‘‘canceled on the right’’ (and hence $F^\#$ on the left)),

(ii) if $F^\# \& F \geq B$, then \mathcal{F} is an epic in Q -Sets (i.e., F is left-cancellable),

(iii) \mathcal{F} is invertible in Q -Sets (of which the inverse is $(F^\#, F)$) iff $A \geq F \& F^\#$ and $F^\# \& F \geq B$.

4. Singletons of Q -sets and complete Q -sets of a quantaloid Q

DEFINITION 4.1. Given a Q -set (X, A) , by a *singleton* S of (X, A) will be meant a pair of matrices, of a ‘‘row’’ $S : 1 \rightarrow X$ and of a ‘‘column’’ $S^\# : X \rightarrow 1$ (with 1 a singleton set) assigning to each $x \in X$ morphisms $s_x : u \rightarrow d(x)$ and $s_x^\# : d(x) \rightarrow u$ of Q , respectively, and having the following properties:

Reproducing Property:

$$s_x = \bigvee_{x' \in X} s_{x'} \& a_{x', x} \quad \text{and} \quad s_x^\# = \bigvee_{x' \in X} a_{x, x'} \& s_{x'}^\#$$

for all $x \in X$, or in matrix terms, $S = S \& A$ and $S^\# = A \& S^\#$;

Singleton Condition: $s_x^\# \& s_{x'} \leq a_{x, x'}$ for all $x, x' \in X$, i.e., $S^\# \& S \leq A$;

Totality: $l \leq \bigvee_{x \in X} s_x \& s_x^\#$ for some idempotent $l : u \rightarrow u$ of Q such that $s_x = l \& s_x$ and $s_x^\# = s_x^\# \& l$ for all $x \in X$, i.e., $\{l\} \leq S \& S^\#$ with $S = \{l\} \& S$, $S^\# = S^\# \& \{l\}$.

We shall sometimes write $d(S)$ for u and shall write \tilde{a}_S for $\bigvee_{x \in X} s_x \& s_x^\#$. It easily verified that, for any strict element $x \in X$ of a Q -set (X, A) , the pair $\mathcal{A}_x = (A_x, A_x^\#)$ consisting of the row $A_x = (a_{x, x'})_{x' \in X}$ and of the column $A_x^\# = (a_{x', x})_{x' \in X}$ of the structural matrix A is a singleton of (X, A) (with $l = a_x$).

PROPOSITION 4.2. Let $\tilde{X} = (\tilde{X}_u)_{u \in Q_0}$ be the family of sets of all singletons of a Q -set (X, A) . Let \tilde{A} be the matrix defined by: $\tilde{a}_{S, T} = S \& T^\#$ for all $S = (S, S^\#)$, $T = (T, T^\#) \in \tilde{X}$. Then the pair (\tilde{X}, \tilde{A}) forms a strict Q -set.

Now we turn to an important class of strict and separated Q -sets of an arbitrary quantaloid Q having the property that each singleton of a Q -set (X, A) is determined by a unique element of X . We need the following analog of Definition 4.15 [3].

DEFINITION 4.3. Let (Y, B) be a Q -set.

(i) We say (X, A) is a *sub- Q -set* of (Y, B) and write $(X, A) \subseteq (Y, B)$ to mean that $X_u \subseteq Y_u$ for each $u \in Q_0$ and the matrix A obtained by restricting B to elements of X makes (X, A) into a Q -set such that

$$b_{x, y} = \bigvee_{x' \in X} a_{x, x'} \& b_{x', y} \quad \text{and} \quad b_{y, x} = \bigvee_{x' \in X} b_{y, x'} \& a_{x', x}$$

for all $x \in X$ and $y \in Y$. (By the way, if (Y, B) is strict, then this condition is always satisfied.)

(ii) We say $(X, A) \subseteq (Y, B)$ generates (Y, B) whenever

$$b_{y,y'} = \bigvee_{x \in X} b_{y,x} \& b_{x,y'}$$

for all $y, y' \in Y$.

Note that henceforth we shall keep the notation $(\underline{X}, \underline{A})$ for the sub- Q -set of a Q -set (X, A) of all strict elements of X .

DEFINITION 4.4. If $(\underline{X}, \underline{A}) \subseteq (X, A)$ generates (X, A) , then the Q -set (X, A) is said to be *strictly generated*.

PROPOSITION 4.5. For a strictly generated Q -set (X, A) , the Q -set (\tilde{X}, \tilde{A}) of all singletons of (X, A) is separated, i.e., if $\tilde{a}_{S', S''} = \tilde{a}_{S'', S''}$ and $\tilde{a}_{S'', S} = \tilde{a}_{S'', S'}$ for all $S'' \in \tilde{X}$, then $S = S'$.

PROPOSITION 4.6. Let $(X, A) \subseteq (Y, B)$ be a sub- Q -set of (Y, B) generating it. For any singleton T of (Y, B) , the restriction ${}_X T$ of T to X is also a singleton of (X, A) satisfying the condition that $\tilde{a}_{{}_X T} = \tilde{b}_T$.

PROPOSITION 4.7. Let $(X, A) \subseteq (Y, B)$ be Q -sets. Then every singleton S of (X, A) extends to a singleton S' of (Y, B) (with $s'_x = s_x$ and $s'^{\#}_x = s^{\#}_x$ for all $x \in X$). Among possible extensions of a singleton $S \in \tilde{X}$ there is the "bottom" extension, the singleton ${}^Y S$ of (Y, B) defined by

$${}^Y s_y = \bigvee_{x \in X} s_x \& b_{x,y} \quad \text{and} \quad {}^Y s_y^{\#} = \bigvee_{x \in X} b_{y,x} \& s_x^{\#}$$

for all $y \in Y$. This singleton has the properties that $\tilde{b}_{{}^Y S} = \tilde{a}_S$ and that ${}^Y S \leq S'$ for any singleton S' of (Y, B) which extends S . The same holds for any pair of such extensions: if ${}^Y S, {}^Y T \in \tilde{Y}$ is a pair of bottom extensions of $S, T \in \tilde{X}$ to singletons of (Y, B) , then $\tilde{b}_{{}^Y S, {}^Y T} = \tilde{a}_{S, T}$, while, for any other extensions $S', T' \in \tilde{Y}$ of $S, T \in \tilde{X}$ (if they exist), $\tilde{b}_{S', T'} \geq \tilde{a}_{S, T}$, in particular, $\tilde{b}_{S'} > \tilde{a}_S$ (for $S' \neq {}^Y S$).

We now come to the key property of the Q -set (\tilde{X}, \tilde{A}) of all singletons of a strictly generated Q -set presented in the next proposition.

PROPOSITION 4.8. Let $\mathcal{K} = (K, K^{\#})$ be a singleton of the Q -set (\tilde{X}, \tilde{A}) of all singletons of a strictly generated Q -set (X, A) . Then there exists a unique element $T \in \tilde{X}$ for which $k_S = \tilde{a}_{T, S}$ and $k^{\#}_S = \tilde{a}_{S, T}$ for all $S \in \tilde{X}$.

This is formalized in the following

DEFINITION 4.9 (cf. Definition 17 [6]). A strict and separated Q -set (X, A) will be said to be complete provided that each singleton $\mathcal{S} = (S, S^\#)$ of (X, A) is of the form: $S = A_x$, i.e., with $S = A_x = (a_{x,x'})_{x' \in X}$ and $S^\# = A^\# = (a_{x',x})_{x' \in X}$, for some (unique) element $x \in X$.

Now we are going to define an adjunction from the full subcategory SGQ -Sets of strictly generated Q -sets of the category Q -Sets to the full subcategory CQ -Sets of complete Q -sets of Q -Sets adapting the construction for quantales considered by C.J. Mulvey and M. Nawaz [6].

PROPOSITION 4.10. Let \sim be the mapping which associates with every strictly generated Q -set (X, A) the complete Q -set (\tilde{X}, \tilde{A}) of all singletons of (X, A) and the morphism mapping which associates with every bimodule $\mathcal{F} : (X, A) \rightarrow (Y, B)$ the bimodule $\tilde{\mathcal{F}} : (\tilde{X}, \tilde{A}) \rightarrow (\tilde{Y}, \tilde{B})$ defined by

$$\tilde{f}_{\mathcal{S}, \mathcal{T}} = \bigvee_{\underline{x} \in X} \bigvee_{\underline{y} \in Y} s_{\underline{x}} \& f_{\underline{x}, \underline{y}} \& t_{\underline{y}}^\# \text{ and } \tilde{f}_{\mathcal{T}, \mathcal{S}} = \bigvee_{\underline{y} \in Y} \bigvee_{\underline{x} \in X} t_{\underline{y}} \& f_{\underline{y}, \underline{x}}^\# \& s_{\underline{x}}^\#.$$

Then \sim is a (covariant) functor from the category SGQ -Sets of strictly generated Q -sets to the category CQ -Sets of complete Q -sets.

Finally, we arrive at

Theorem 4.11 (cf. Corollary 19 [6]). *The functors*

$$SGQ\text{-Sets} \overset{\sim}{\cong} CQ\text{-Sets}$$

establish an equivalence of categories, where I is the embedding of the subcategory CQ -Sets of complete Q -sets in the category SGQ -Sets of strictly generated Q -sets.

5. Presheaves on a quantaloid Q

Henceforth we shall solely work with strict Q -sets.

DEFINITION 5.1. By a presheaf on Q will be meant a strict Q -set (X, A) together with *restriction*: a partial mapping $[\] : Q_1 \times X \times Q_1 \rightarrow X$ (more precisely, a "matrix" $[\] = ([\]_{u,v})_{\substack{u \in Q_0 \\ v \in Q_0}}$ of partial mappings $[\]_{u,v} : Q(u, v) \times X_v \times Q(v, u) \rightarrow X_u$) from the triplets $(p, x, p^\#) \in Q_1 \times X \times Q_1$ with $dom(p) = cod(p^\#)$ and $cod(p) = dom(p^\#) = d(x)$ (keeping $d(p, x, p^\#) = dom(p)$) such that $a_x \& p^\# \& p \& a_x \leq a_x$, $p \& a_x \leq p \& a_x \& p^\# \& p \& a_x$, and $a_x \& p^\# \leq a_x \& p^\# \& p \& a_x \& p^\#$ (the second and the third of which are actually equalities owing to the first condition), which will be referred to as *restrictable triplets* of $Q_1 \times X \times Q_1$ to the elements $p[x]p^\#$ of X satisfying the compatibility conditions that: $q[(p[x]p^\#)]q^\# =$

$(q \& p)[x](p^\# \& q^\#)$, $a_x[x]a_x = x$, and $a_{p[x]p^\#, p'[x']p'^\#} = p \& a_{x,x'} \& p'^\#$ for all restrictable triplets $(p, x, p^\#)$, $(p', x, p'^\#)$, and $(q, p[x]p^\#, q^\#)$ of $Q_1 \times X \times Q_1$. A presheaf $(X, A, [])$ on Q is *separated* whenever the underlying Q -set (X, A) is so.

DEFINITION 5.2 (cf. [6]). By the *canonical presheaf* $(X, A, [])$ on Q determined by a complete Q -set (X, A) will be meant the presheaf of which the underlying Q -set is (X, A) itself and of which the restriction is uniquely determined by requiring that $a_{p[x]p^\#, x''} = p \& a_{x,x''}$ and $a_{x'', p[x]p^\#} = a_{x'', x} \& p^\#$ for any restrictable triplet $(p, x, p^\#)$ of $Q_1 \times X \times Q_1$ and for all elements $x'' \in X$.

DEFINITION 5.3 (cf. Definition 23 [6]). By a *map of presheaves* $f : (X, A, []) \rightarrow (Y, B, [])$ on Q will be meant a map $f : X \rightarrow Y$ in $Sets/Q_0$ satisfying the following:

Strictness: $d(x) = d(fx)$, $b_{f x, y} = a_x \& b_{f x, y}$, and $b_{y, f x} = b_{y, f x} \& a_x$ for all $x \in X$ and $y \in Y$;

Isotonicity: $a_{x, x'} \leq b_{f x, f x'}$ for all $x, x' \in X$;

Preservation of Restriction: if $(p, x, p^\#)$ is restrictable, then $(p, f x, p^\#)$ is also restrictable and $f(p[x]p^\#) = p[f x]p^\#$.

DEFINITION 5.4 (cf. Definition 24 [6]). By the *canonical map of presheaves*

$$f^\Gamma : (X, A, []) \rightarrow (Y, B, [])$$

determined by a bimodule $F : (X, A) \rightarrow (Y, B)$ between complete Q -sets will be meant the map $f^\Gamma : X \rightarrow Y$ in $Sets/Q_0$, every value $f^\Gamma x$ of which is uniquely determined by requiring that $b_{f^\Gamma x, y} = f_{x, y}$ and $b_{y, f^\Gamma x} = f_{y, x}^\#$ for all $y \in Y$.

For any quantaloid Q , presheaves on Q together with maps of presheaves on Q form a category, which we shall denote by $Q - Psh$. Moreover, the assignment $\mathcal{F} \mapsto f^\Gamma$ determines a functor from the category $CQ - Sets$ of complete Q -sets to the category $Q - Psh$ of presheaves on Q , which we shall denote by Γ . We are going to establish the existence of the adjoint to Γ . First, we present several results from [2] (more precisely, their "presheaf" versions).

PROPOSITION 5.5 (Proposition 2.7 [2]). Every map $f : (X, A, []) \rightarrow (Y, B, [])$ of presheaves on Q determines a bimodule $\mathcal{F}^U = (F^U, F^{U\#})$ from (X, A) to (Y, B) by the relations $f_{x, y}^U = b_{f x, y}$ and $f_{y, x}^{U\#} = b_{y, f x}$ for all $x \in X$ and $y \in Y$.

PROPOSITION 5.6 (Proposition 2.9 [2]). The assignment $f \mapsto \mathcal{F}^U$ determines a functor from the category $Q - Psh$ of presheaves on Q to the full subcategory $SQ - Sets$ of strict Q -sets of the category $Q - Sets$ (or of the category $SGQ - Sets$), which we shall denote by $U : Q - Psh \rightarrow SQ - Sets$.

It is clear that the composite ΓU (rightwards) is none other than the inclusion functor $I : CQ - Sets \rightarrow SQ - Sets$, since the assignment $\mathcal{F} \mapsto f^\Gamma$ is one-to-one. The following proposition generalize Theorem 25 [6].

Theorem 5.7. For any quantaloid Q , the functors

$$Q\text{-Psh} \xrightleftharpoons[\Gamma]{\Sigma=U\sim} CQ\text{-Sets}$$

are adjoint, where \sim is the functor from $SQ\text{-Sets}$ to $CQ\text{-Sets}$ (introduced in Proposition 4.10).

6. Sheaves on a quantaloid

In this section we describe a novel "sheaf condition" on a separated presheaf, corresponding to that of the completeness of a Q -set.

DEFINITION 6.1. We say that the separated presheaf $(X, A, [\])$ on Q is the sheaf on Q if it satisfies *sheaf condition*:

(i) for every singleton \mathcal{S} of the underlying Q -set (X, A) , there exist "enough" restrictable triplets $(s_x, x, s_x^\#)$ in the sense that

$$s_x = \bigvee \{s_{x'} \& a_{x',x} \mid x' \in X, (s_{x'}, x', s_{x'}^\#) \text{ restrictable}\}$$

and

$$s_x^\# = \bigvee \{a_{x,x'} \& s_{x'}^\# \mid x' \in X, (s_{x'}, x', s_{x'}^\#) \text{ restrictable}\}$$

for all $x \in X$ (noting that $(s_x, x, s_x^\#)$ is restrictable iff $s_x = s_x \& s_x^\# \& s_x$ and $s_x^\# = s_x^\# \& s_x \& s_x^\#$);

(ii) if the subset $J \subseteq X_u$ (for some $u \in Q_0$) is such that the pair $\mathcal{E} = (E, E^\#)$ (with $E = (e_x)_{x \in J}$, $E^\# = (e_x^\#)_{x \in J}$, and $e_x = e_x^\# = a_x$ ("diagonal" element of A) for $x \in J$) constitutes a singleton of the sub- Q -set $(J, J A) \subseteq (X, A)$, then its "bottom" extension ${}^X \mathcal{E} = ({}^X E, {}^X E^\#)$ to a singleton of (X, A) obtained by (see Proposition 4.7):

$${}^X e_{x'} = \bigvee_{x \in J} e_x \& a_{x,x'} (= \bigvee_{x \in J} a_{x,x'}) \text{ and } {}^X e_{x'}^\# = \bigvee_{x \in J} a_{x',x} \& e_x^\# (= \bigvee_{x \in J} a_{x',x})$$

for all $x' \in X$, is of the form:

$${}^X \mathcal{E} = \mathcal{A}_y$$

for some (unique) element $y \in X$ (where $\mathcal{A}_y = ((a_{y,x})_{x \in X}, (a_{x,y})_{x \in X})$).

PROPOSITION 6.2. The canonical presheaf $(X, A, [\])$ determined by any complete Q -set is a sheaf.

With these remarks, denoting for a quantaloid Q by $Q - Sh$ the full subcategory of the category $Q - Psh$ of presheaves on Q determined by sheaves on Q , we see that the functor $\Gamma : CQ - Sets \rightarrow Q - Psh$ (determined in Definition 5.4) that assigns to each complete Q -set its canonical presheaf may in fact be considered as a functor into the category of sheaves on Q , and that the functor $U : Q - Psh \rightarrow SQ - Sets$ (introduced in Proposition 5.6 that assigns to each presheaf its underlying Q -set) restricted to $Q - Sh$ may be regarded as a functor into $CQ - Sets$. We have the following analog of Theorem 29 [6], and also of Theorem 4.13 [3].

Theorem 6.3. *For a quantaloid Q , the categories of complete Q -sets and of sheaves on Q are isomorphic by the functors*

$$CQ - Sets \begin{array}{c} \xrightarrow{\Gamma} \\ \xleftarrow{U} \end{array} Q - Sh$$

that assign respectively to each complete Q -set its canonical presheaf, and to each sheaf its underlying Q -set.

References

- B.I. F. Borceux, G. van den Bossche, Quantales and their sheaves, *Order* 3, 61–87 (1986).
 G. van den Bossche, Quantaloids and non-commutative ring representations, *Appl. Categ. Structures* 3, 305–320 (1995).
 F.M.P. Fourman, D.S. Scott, Sheaves and logic, in: *Applications of Sheaves, Lecture Notes in Mathematics* 753, Springer-Verlag, Berlin, Heidelberg, New York (1979) pp. 302–401.
 U. Höhle, M -valued sets and sheaves over integral cl -monoids, in: *Applications of Category Theory to Fuzzy Subsets*, eds. S.E. Rodabaugh et al., Kluwer, Boston (1992), pp. 33–72.
 U. Höhle, GL -Quantales: Q -valued sets and their singletons, *Studia Logica* 61, 123–148 (1998).
 C.J. Mulvey, M. Nawaz, Quantales: quantal sets, in: *Non-Classical Logics and their Applications to Fuzzy Subsets*, eds. U. Höhle, E.P. Klement, Kluwer, Boston (1995), pp. 159–217.
 K.I. Rosenthal, *Quantales and Their Applications, Pitman Research Notes in Mathematics* 234, Longman, Burnt Mill, Harlow (1990).
 K.I. Rosenthal, *The Theory of Quantaloids*, Longman, Harlow (1996).

Pluoštai virš kvantaloidų

R.P. Gyls

Tiriamos aiškus su reikšmėmis iš kvantaloido. Išvestos formulės apibendrina nesenus C.J. Mulvey ir M. Nawaz rezultatus.