





# Existence and approximate controllability results for time-fractional stochastic Navier–Stokes equations

Renu Chaudhary<sup>a,b,1</sup> , Simeon Reich<sup>a,2</sup>, Juan J. Nieto<sup>c,3</sup> 

<sup>a</sup>Department of Mathematics,  
The Technion – Israel Institute of Technology,  
32000 Haifa, Israel  
[renu.chaudhary@thws.de](mailto:renu.chaudhary@thws.de)

<sup>b</sup>Faculty of Applied Natural Sciences and Humanities (FANG),  
Technical University of Applied Sciences Würzburg-Schweinfurt,  
97421 Schweinfurt, Germany

<sup>c</sup>CITMAga, Departamento de Estatística,  
Análise Matemática e Optimización Universidade de Santiago de Compostela,  
15782 Santiago de Compostela, Spain

**Received:** June 29, 2025 / **Revised:** January 16, 2026 / **Published online:** April 21, 2026

**Abstract.** This paper deals with time-fractional stochastic Navier–Stokes equations, which are characterized by the coexistence of stochastic noise and a fractional power of the Laplacian. We establish sufficient conditions for the existence and approximate controllability of a unique mild solution to time-fractional stochastic Navier–Stokes equations. Using a fixed point technique, we first demonstrate the existence and uniqueness of a mild solution to the equation under consideration. We then establish approximate controllability results by using the concepts of fractional calculus, semigroup theory, functional analysis, and stochastic analysis.

**Keywords:** approximate controllability, time-fractional Navier–Stokes equations, stochastic analysis, semigroup theory, fixed point technique.

## 1 Introduction

The origin of the theory of stochastic Navier–Stokes equations (SNSEs, for short) dates back to the work of Landau and Lifshitz [20] in their 1959 book entitled “Fluid Mechanics”. Later on, after the pioneering work of Bensoussan and Temam [2] on the

---

<sup>1</sup>Corresponding author.

<sup>2</sup>The author was supported by the Israel Science Foundation (grant No. 820/17), the Fund for the Promotion of Research at the Technion (grant No. 2001893), and the Technion General Research Fund (grant No. 2016723).

<sup>3</sup>The author was supported by the Agencia Estatal de Investigación (Spain) under grant PID2020-113275GB-I00, funded by MCIN/AEI/10.13039/501100011033, and by the European Regional Development Fund (ERDF “A way of making Europe”), as well as by the Xunta de Galicia under grant ED431C 2019/02 (Competitive Reference Research Groups, 2019–2022).

mathematical viewpoint of SNSEs, these equations have been widely studied and applied to characterize several phenomena in various domains, especially in the fields of fluid dynamics, gas dynamics, nonlinear acoustics, and traffic flow. The stochasticity introduced in the well-known Navier–Stokes equation might help us comprehend physical processes and the mechanisms of fluid turbulence in a more sophisticated manner. The SNSEs provide a relatively simple method for investigating turbulence, distortion caused by laminar momentum transport, and the decay of dissipation layers formed as a result. In the last few years, significant progress has been made in the mathematical and numerical analysis of SNSEs. In particular, Da Prato and Debussche [6] and Brzeźniak [3] studied two-dimensional SNSEs. Kim [17] established the existence of strong solutions to SNSEs in  $\mathbb{R}^3$ . Recently, Hofmanova [13] has derived global existence and non-uniqueness results for three-dimensional SNSEs.

It is well known that fractional derivatives (rather than integer-order derivatives) have proven to be highly useful in analyzing many real-world problems because of their ability to represent long memory processes; see [16, 22, 24]. Experiments show that fractional derivatives are among of the top tools for modeling anomalous diffusion processes and viscoelasticity; see, for instance, [4, 10, 11, 26]. Consequently, fractional-order Navier–Stokes equations (FNSEs, for short) can be used to model more precisely anomalous diffusion in fractal media. As a result, numerous researchers have conducted in-depth studies regarding FNSEs. For example, de Carvalho-Neto and Planas [8] established the existence of mild solution to FNSEs in  $\mathbb{R}^n$ ; Momani and Odibat [23] studied analytical solutions of FNSEs using different methods; Zhou and Peng [29] established the existence and uniqueness of local and global mild solutions to FNSEs. Xu et al. [28] and Zou et al. [31] have analyzed FNSEs driven by fractional Brownian motion. Recently, Han [12] studied  $L_p$ -solvability for stochastic time fractional Burgers' equations driven by multiplicative space-time white noise. The author established conditions for the existence and uniqueness and Hölder regularity of the local and global solution.

Controllability plays an essential role in the design and analysis of control systems. Starting from the seminal work of Kalman [14] in 1963, the notion of controllability helps one to understand whether a particular dynamical control system can be controlled or not, and if it can, what the optimum control should be in order to obtain the intended outcome. Our understanding of the controllability of fractional control systems has greatly increased in recent years as more scientists have become interested in the topic of controllability and its applications in biological science, economics, aerospace engineering, and electrical engineering; see, for instance, Barnett [1], Curtain and Zwart [5], Dauer and Mahmudov [7], Karthikeyan et al. [15], Sakthivel et al. [25], and references therein. In spite of its being an important topic, only a few papers deal with the controllability of FNSEs. Xi et al. [27] established approximate controllability results for time-fractional Navier–Stokes equations involving time delay, and Liao et al. [21] have recently presented global controllability results for Navier–Stokes equations. However, the existing literature does not address the question of existence and approximate controllability of FNSEs involving randomness and fractional powers of the Laplacian. Therefore it is quite natural and interesting to establish existence and controllability results for time-fractional SNSEs.

This paper makes significant contributions to this unexplored domain by investigating the existence and approximate controllability of solutions to time-fractional SNSEs. We develop an innovative application of fixed point theorems tailored to handle the intricacies of fractional stochastic systems, establishing the existence and uniqueness of mild solutions under conditions not previously addressed. By examining the approximate controllability within this complex setting, we provide new theoretical results that expand on existing controllability concepts, accounting for both fractional and random influences. Our results have practical implications for modeling and controlling fluid dynamics phenomena where anomalous diffusion and random effects are prevalent, such as pollutant dispersion in atmospheric sciences and intricately controlled flows in biotechnology. We mainly focus on the following time-fractional SNSE. This SNSE is the simplest representative equation of the tri-interaction between wave steepening, small dissipation and random perturbations, which are represented as the nonlinearity, the fractional power of the Laplacian and the stochastic process, respectively, in a bounded domain  $\mathcal{O} \subset \mathbb{R}^d$  ( $1 \leq d \leq 3$ ) with a smooth boundary  $\partial\mathcal{O}$ :

$$\begin{aligned} \partial_t^\eta z(t, x) + \nu(-\Delta)^{\alpha/2} z(t, x) - (z(t, x) \cdot \nabla) z(t, x) - \nabla \rho(t, x) \\ = Cv(t, x) + \tilde{h}(t, z(t, x)) \frac{dW(t)}{dt}, \end{aligned} \quad (1)$$

where  $(t, x) \in (0, T] \times \mathcal{O}$ , with the incompressibility condition

$$\nabla \cdot z(t, x) = 0, \quad (t, x) \in (0, T] \times \mathcal{O}, \quad (2)$$

the Dirichlet boundary condition


$$z(t, x) = 0, \quad (t, x) \in (0, T] \times \partial\mathcal{O}, \quad (3)$$

and the initial conditions

$$z(0, x) = z_0(x), \quad x \in \mathcal{O}. \quad (4)$$

Here  $\eta \in (0, 1)$ ,  $z(t, x)$  denotes the velocity field at a point  $x \in \mathbb{R}^d$ ,  $\nu > 0$  is the viscosity coefficient,  $\rho$  denotes the associated pressure field,  $C$  represents a linear operator,  $v$  denotes the control function,  $\tilde{h}$  is a nonlinear function, which represents the external force,  $W(t)$  denotes a Wiener process, and the operator  $(-\Delta)^{\alpha/2}$ ,  $\alpha \in (1, 2)$ , denotes a fractional power of the Laplacian; see [9, 19].

The time-fractional stochastic Navier–Stokes equations formulated in this work serve as a robust framework for modeling fluid behavior in scenarios where classical assumptions of instantaneous response and deterministic evolution no longer hold. The fractional time derivative effectively captures memory-dependent dynamics, which are characteristic of viscoelastic or anomalously diffusing fluids, while the stochastic perturbations reflect uncertainties and fluctuations inherent in many natural and engineered systems. Such a model becomes particularly relevant in applications ranging from subsurface contaminant transport and atmospheric dispersion of pollutants to microfluidic flows in biomedical devices. In these settings, the combination of long-range temporal correlations



**Figure 1.** Logical structure of the paper: lemmas, definitions, assumptions, and theorems with explicit dependencies.

and random influences plays a crucial role, and understanding the controllability of such systems can significantly inform the design of efficient monitoring, control, and optimization strategies.

## 2 Notations and preliminaries

For  $1 < p < \infty$ ,  $L^p(\mathcal{O})$  denotes the Lebesgue space, and  $W^{k,p}(\mathcal{O})$  denotes the Sobolev spaces with  $H^k(\mathcal{O}) := W^{k,2}(\mathcal{O})$ . For  $p = 2$ ,  $L^2(\mathcal{O}) =: H$  denotes the Hilbert space with inner product  $\langle \cdot, \cdot \rangle$ . Let  $C^\infty(\mathcal{O})$  denote the space of all infinitely differentiable functions and  $C_0^\infty(\mathcal{O}) := \{z \in (C^\infty(\mathcal{O}))^d : \nabla \cdot z = 0, z \text{ has compact support in } \mathcal{O}\}$ . Let  $H_0^1(\mathcal{O})$  denote the closure of  $C_0^\infty(\mathcal{O})$  in  $(L^p(\mathcal{O}))^d$ .

Let  $(\Sigma, \mu, \{\mu_t\}_{t \geq 0}, \mathbf{P})$  be a filtered complete probability space with normal filtration  $\{\mu_t\}_{t \geq 0}$ . The Wiener process  $\{W(t), t \geq 0\}$  possesses a linear, bounded covariance operator  $Q \geq 0$  of finite trace, with  $\text{Tr}(Q) = \sum_{m=1}^\infty \nu_m = \nu < \infty$  and  $Qc_m = \nu_m c_m$ , where  $\{c_m, m \in \mathbb{N}\}$  denotes a complete orthonormal basis for  $H$ . If  $\{\omega_m\}_{m \in \mathbb{N}}$  is a sequence of one-dimensional Wiener processes, then

$$W(t) = \sum_{m=1}^\infty \sqrt{\nu_m} \omega_m(t) c_m, \quad t \geq 0.$$

$L_0^2 = L^2(Q^{1/2}(U), H)$  denotes the Hilbert space of Hilbert–Schmidt operators from  $Q^{1/2}(U)$  to  $H$  endowed with the norm

$$\|\psi\|_{L_0^2} = \left( \sum_{m=1}^\infty \|\psi \nu_m\|^2 \right)^{1/2}, \quad \psi \in L_0^2.$$

Furthermore, let  $L^p(\Sigma; H)$  be the Hilbert space of  $H$ -valued random variables with norm

$$\|z(\cdot)\|_{L^p(\Sigma; H)} = (\mathbf{E} \|z(\cdot)\|_H^p)^{1/p},$$

where  $\mathbf{E}$  represents the expectation with respect to the measure  $\mathbf{P}$  defined by

$$\mathbf{E}\|z(\cdot)\|_H^p := \int_{\Sigma} \|z(\omega)\|_H^p d\mathbf{P}(\omega) < \infty, \quad \omega \in \Sigma.$$

Let  $P$  be the Helmholtz projection operator defined on  $(L^p(\mathcal{O}))^d$  with range  $H_0^1(\mathcal{O})$ . Let  $A = -\nu P\Delta$  denote the Stokes operator in  $H_0^1(\mathcal{O})$  with domain  $D(A) = \{z \in H_0^1(\mathcal{O}) \cap H^2(\mathcal{O}): z(t, x) = 0 \forall (t, x) \in (0, T] \times \partial\mathcal{O}\}$ . Since  $\mathcal{O}$  is bounded, the inverse  $A^{-1}$  exists and is a compact operator on  $H$ . Furthermore, the fractional powers of  $A$ , that is,  $A^{\alpha/2} = A_\alpha = \nu P(-\Delta)^{\alpha/2}$ , can be defined by

$$A^{\alpha/2}v_m = e_m^{\alpha/2}v_m$$

with domain

$$\mathcal{H}^\alpha = D(A^{\alpha/2}) = \left\{ z \in H: \|z\|_{\mathcal{H}^\alpha}^2 = \sum_{m=1}^\infty e_m^{\alpha/2}u_m^2 < \infty \right\},$$

where  $u_m = \langle z, v_m \rangle$ . We set  $\|z\|_{\mathcal{H}^\alpha} := \|A^{\alpha/2}z\|$  and consider the associated dual space  $\mathcal{H}^{-\alpha}$  with the inverse operator  $A^{-\alpha/2}$ . The operator  $-A_\alpha$  generates an analytic semigroup  $S_\alpha(t) = e^{-tA_\alpha}$  of operators, which are compact for  $t > 0$ . The control function  $v \in L_\mu^p([0, T], U)$ , where  $U$  is the separable Hilbert space. For simplicity of notation, the set of admissible controls is denoted by  $U_{\text{ad}} = L_\mu^p([0, T], U)$ . The mapping  $C : U \rightarrow H$  is a bounded linear operator. We also consider the bilinear operator  $G(z, w) := -P(z \cdot \nabla)w$  with  $D(G) = H_0^1(\mathcal{O})$ . In a slight abuse of notation, we write  $G(z) := G(z, z)$ .

Applying the Helmholtz–Hodge projection operator  $P$  to the time-fractional SNSE (1) subject to conditions (2)–(4), we obtain the following abstract equation:

$$\begin{aligned} {}^C D^\eta z(t) &= -A_\alpha z(t) + G(z(t)) + Cv(t) + \hbar(t, z(t)) \frac{dW(t)}{dt}, \quad t \in (0, T], \\ z(0) &= z_0, \end{aligned} \tag{5}$$

where  ${}^C D^\eta$  denotes the Caputo fractional derivative of order  $\eta \in (0, 1)$ . Here  $-A_\alpha$  is the infinitesimal generator of an analytic semigroup  $\{S_\alpha(t), t \geq 0\}$ . The initial value  $z_0$  is an  $H^\alpha$ -valued  $\mu_0$ -measurable random variable, which is independent of  $W$ . Instead of  $Pv(t)$  and  $P\hbar(t, z(t))$ , we will use the notation  $v(t)$  and  $\hbar(t, z(t))$ , respectively.

**Definition 1.** (See [24].) The Caputo fractional derivative of  $z \in C^m([0, T])$  of order  $\eta$ ,  $m - 1 < \eta \leq m$ , is defined by

$${}^C D_{0+}^\eta z(t) := \frac{1}{\Gamma(m - \eta)} \int_0^t (t - r)^{m-\eta-1} \frac{d^m}{ds^m} z(r) dr,$$

where  $z^{(m-1)}$  is absolutely continuous in every compact interval  $[0, T]$ ,  $T > 0$ .

**Lemma 1.** (See [30].) For any  $\alpha > 0$ , the operator  $-A_\alpha$  generates an analytic semigroup  $S_\alpha(t) = e^{-tA_\alpha}$ ,  $t \geq 0$ , on  $L^p$ . Moreover, we have

$$\|A_\beta S_\alpha(t)\|_{\mathcal{L}(L^p)} \leq C_{\alpha,\beta} t^{-\beta/\alpha}, \quad t > 0,$$

where  $\beta \geq 0$ , the constant  $C_{\alpha,\beta} > 0$  depends on  $\alpha$  and  $\beta$ , and  $\mathcal{L}(L^p)$  denotes the Banach space of all bounded linear operators from  $L^p$  into itself.

**Lemma 2.** (See [18].) For any  $p \geq 2$ ,  $0 \leq s_1 < s_2 \leq T$ , and predictable stochastic process  $\chi : [0, T] \times \Sigma \rightarrow L^2_0$  such that

$$\mathbf{E} \left( \int_0^T \|\chi(s)\|_{L^2_0}^2 ds \right)^{p/2} < \infty,$$

we have

$$\mathbf{E} \left\| \int_{s_1}^{s_2} \chi(s) dW(s) \right\|^p \leq \kappa(p) \mathbf{E} \left( \int_{s_1}^{s_2} \|\chi(s)\|_{L^2_0}^2 ds \right)^{p/2}.$$

Here the constant

$$\kappa(p) = \left( \frac{p}{2}(p-1) \right)^{p/2} \left( \frac{p}{p-1} \right)^{p(p/2-1)}.$$

**Definition 2.** (See [30].) A  $\mu_t$ -adapted stochastic process  $\{z(t)\}_{t \in [0, T]}$  is said to be a mild solution of (5) if for each  $v \in U_{ad}$ ,  $\{z(t)\}_{t \in [0, T]} \in C([0, T], \mathcal{H}^\alpha)$   $\mathbf{P}$ -a.s. and

$$\begin{aligned} z(t) &= M_\eta(t)z_0 + \int_0^t (t-r)^{\eta-1} M_{\eta,\eta}(t-r) [G(z(r)) + Cv(r)] dr \\ &+ \int_0^t (t-r)^{\eta-1} M_{\eta,\eta}(t-r) h(r, z(r)) dW(r). \end{aligned} \tag{6}$$

Here  $M_\eta(t)$  and  $M_{\eta,\eta}(t)$  denote the generalized Mittag-Leffler operators given by

$$M_\eta(t) = \int_0^\infty K_\eta(s) S_\alpha(t^\eta s) ds \quad \text{and} \quad M_{\eta,\eta}(t) = \int_0^\infty \eta s K_\eta(s) S_\alpha(t^\eta s) ds,$$

where  $K_\eta : \mathbb{C} \rightarrow \mathbb{C}$  is the Mainardi function, which is defined by

$$K_\eta(z) := \sum_{m=0}^\infty \frac{(-1)^m (z)^m}{m! \Gamma(-\eta m + 1 - \eta)}$$

for each  $\eta \in (0, 1)$ .

Next, we recall some properties of  $M_\eta(t)$  and  $M_{\eta,\eta}(t)$ , which are demonstrated in [30].

**Lemma 3.** (See [30].) *The operators  $M_\eta(t)$  and  $M_{\eta,\eta}(t)$ ,  $t > 0$ , are linear bounded operators such that for  $0 \leq \beta < \alpha < 2$ , we have*

$$\|M_\eta(t)z\|_{\mathcal{H}^\beta} \leq C_\alpha t^{-\eta\beta/\alpha} \|z\| \quad \text{and} \quad \|M_{\eta,\eta}(t)z\|_{\mathcal{H}^\beta} \leq C_\eta t^{-\eta\beta/\alpha} \|z\|,$$

where

$$C_\alpha = \frac{C_{\alpha,\beta}\Gamma(1 - \frac{\beta}{\alpha})}{\Gamma(1 - \frac{\eta\beta}{\alpha})} \quad \text{and} \quad C_\eta = \frac{\eta C_{\alpha,\beta}\Gamma(2 - \frac{\beta}{\alpha})}{\Gamma(1 + \eta(1 - \frac{\beta}{\alpha}))}.$$

**Lemma 4.** (See [30].) *The operators  $M_\eta(t)$  and  $M_{\eta,\eta}(t)$ ,  $t > 0$ , are strongly continuous, and for any  $0 < T_0 \leq \tau_1 < \tau_2 \leq T$  and  $0 \leq \beta < \alpha < 2$ , we have*

$$\|(M_\eta(\tau_2) - M_\eta(\tau_1))z\|_{\mathcal{H}^\beta} \leq C_{\alpha\beta}(\tau_2 - \tau_1)^{\eta\beta/\alpha} \|z\|$$

and

$$\|(M_{\eta,\eta}(\tau_2) - M_{\eta,\eta}(\tau_1))z\|_{\mathcal{H}^\beta} \leq C_{\eta\beta}(\tau_2 - \tau_1)^{\eta\beta/\alpha} \|z\|,$$

where

$$C_{\alpha\beta} = \frac{\alpha C_{\alpha,\beta}\Gamma(1 - \frac{\beta}{\alpha})}{\beta T_0^{2\eta\beta/\alpha}\Gamma(1 - \frac{\eta\beta}{\alpha})} \quad \text{and} \quad C_{\eta\beta} = \frac{\alpha\eta C_{\alpha,\beta}\Gamma(2 - \frac{\beta}{\alpha})}{\beta T_0^{2\eta\beta/\alpha}\Gamma(1 + \eta(1 - \frac{\beta}{\alpha}))}.$$

Now we define the stochastic controllability operator  $L_T \in \mathcal{L}(U_{\text{ad}}, L^p(\Sigma, H))$  by

$$L_T v := \int_0^T (T-s)^{\eta-1} M_{\eta,\eta}(T-s) C v(s) ds$$

and the corresponding adjoint operator  $L_T^* : L^p(\Sigma, H) \rightarrow U_{\text{ad}}$  by

$$L_T^* z = C^* M_{\eta,\eta}^*(T-s) \mathbf{E}\{z \mid \mu_t\}.$$

Similarly to the Grammian matrix, we have the stochastic Grammian operator

$$\begin{aligned} \mathcal{Y}_0^T &:= L_T(L_T^*)z \\ &= \int_0^T (T-s)^{\eta-1} M_{\eta,\eta}(T-s) C C^* M_{\eta,\eta}^*(T-s) \mathbf{E}\{z \mid \mu_s\} ds. \end{aligned}$$

Next, we define the reachable set  $K(T) := \{z(T, z_0, v) : v(\cdot) \in U_{\text{ad}}\}$  of (5), which is the set of all final states  $z$  with initial state  $z_0$  and control  $v$  at the terminal time  $T$ .

**Definition 3.** The time-fractional SNSE (5) is called approximately controllable on  $[0, T]$  if  $\overline{K(T)} = L^p(\Sigma, H)$ .

**Lemma 5.** (See [7].) *For any  $z_T \in L^p(\Sigma, H)$ , there exists  $\varphi \in L^p_\mu(\Sigma, L^p((0, T), L^p_0))$  such that  $z_T = E z_T + \int_0^T \varphi(s) dW(s)$ .*

### 3 Existence and controllability results

Assume the following conditions:

- (i) The bounded bilinear operator  $G : H \rightarrow H^{-1}(\mathcal{O})$  satisfies the conditions

$$\|G(z)\|_{H^{-1}} \leq C_1 \|z\|^2$$

and

$$\|G(z) - G(w)\|_{H^{-1}} \leq C_2 (\|z\| + \|w\|) \|z - w\|,$$

where  $C_1$  and  $C_2$  are positive constants.

- (ii)  $\hbar : [0, T] \times H \rightarrow L_0^2$  is a measurable function, which satisfies the conditions

$$\|\hbar(t, z(t))\|_{L_0^2} \leq L_1 (1 + \|z\|)$$

and

$$\|\hbar(t, z(t)) - \hbar(t, w(t))\|_{L_0^2} \leq L_2 (\|z - w\|),$$

where  $L_1$  and  $L_2$  are positive constants.

- (iii) The linear deterministic system corresponding to (5),

$$\begin{aligned} {}^C D^\eta z(t) &= -A_\alpha z(t) + G(z(t)) + Cv(t), \quad t \in (0, T], \\ z(0) &= z_0, \end{aligned}$$

is approximately controllable on  $[t, T]$ , that is, for each  $t \in [0, T]$ , the operator  $\lambda(\lambda I + \Upsilon_0^T) \rightarrow 0$  as  $\lambda \rightarrow 0+$  in the strong operator topology.

For any  $z_T \in L^p(\Sigma, H^\beta)$ , we define the control function

$$\begin{aligned} v^\lambda(t, z) &= C^* M_{\eta, \eta}^*(T - t) \left[ (\lambda I + \Upsilon_0^T)^{-1} (\mathbf{E} z_T - M_\eta(T) z_0) \right. \\ &\quad + \int_0^t (\lambda I + \Upsilon_0^T)^{-1} \varphi(r) dW(r) - \int_0^t (\lambda I + \Upsilon_0^T)^{-1} M_{\eta, \eta}(T - r) G(z(r)) dr \\ &\quad \left. - \int_0^t (\lambda I + \Upsilon_0^T)^{-1} M_{\eta, \eta}(T - r) \hbar(r, z(r)) dW(r) \right]. \end{aligned}$$

**Lemma 6.** For  $p \geq 2$ ,  $0 \leq \beta < \alpha < 2$ , and for all  $z, w \in L^p(\Sigma, H^\beta)$ , we have

$$\begin{aligned} \mathbf{E} \|v^\lambda(t, z) - v^\lambda(t, w)\|^p &\leq \frac{C_v}{\lambda^p} \int_0^t \mathbf{E} \|z(r) - w(r)\|^p dr, \\ \mathbf{E} \|v^\lambda(t, z)\|^p &\leq \frac{C_v}{\lambda^p} \left( 1 + \int_0^t \mathbf{E} \|z(r)\|^p dr \right), \end{aligned} \tag{7}$$

where  $C_v$  denotes a constant.



*Proof.* For  $z, w \in L^p(\Sigma, H^\beta)$ , using the bounds of the linear operator  $M_{\eta,\eta}$  from Lemma 3, Hölder’s inequality, and conditions (i)–(ii), we obtain

$$\begin{aligned}
 & \mathbf{E} \|v^\lambda(t, z) - v^\lambda(t, w)\|^p \\
 & \leq 2^{p-1} \left[ \mathbf{E} \left\| C^* M_{\eta,\eta}^*(T-t) \int_0^t (\lambda I + \mathcal{Y}_0^T)^{-1} M_{\eta,\eta}(T-r) [G(z(r)) - G(w(r))] \, dr \right\|^p \right. \\
 & \quad \left. + \mathbf{E} \left\| C^* M_{\eta,\eta}^*(T-t) \int_0^t (\lambda I + \mathcal{Y}_0^T)^{-1} M_{\eta,\eta}(T-r) \right. \right. \\
 & \quad \quad \left. \left. \times [\tilde{h}(r, z(r)) - \tilde{h}(r, w(r))] \, dW(r) \right\|^p \right] \\
 & \leq 2^{p-1} \|C^*\|^p \|M_{\eta,\eta}^*(T-t)\|^p \\
 & \quad \times \left[ \mathbf{E} \left\| \int_0^t (\lambda I + \mathcal{Y}_0^T)^{-1} A_1 M_{\eta,\eta}(T-r) A_{-1} [G(z(r)) - G(w(r))] \, dr \right\|^p \right. \\
 & \quad \left. + \mathbf{E} \left\| \int_0^t (\lambda I + \mathcal{Y}_0^T)^{-1} M_{\eta,\eta}(T-r) [\tilde{h}(r, z(r)) - \tilde{h}(r, w(r))] \, dW(r) \right\|^p \right] \\
 & \leq 2^{p-1} \frac{\|C^*\|^p}{\lambda^p} C_\eta^p \left[ \mathbf{E} \left\| \int_0^t A_1 M_{\eta,\eta}(T-r) A_{-1} [G(z(r)) - G(w(r))] \, dr \right\|^p \right. \\
 & \quad \left. + \mathbf{E} \left\| \int_0^t M_{\eta,\eta}(T-r) [\tilde{h}(r, z(r)) - \tilde{h}(r, w(r))] \, dW(r) \right\|^p \right] \\
 & \leq 2^{p-1} \frac{\|C^*\|^p}{\lambda^p} C_\eta^p [I_1 + I_2], \tag{8}
 \end{aligned}$$

where

$$I_1 = \mathbf{E} \left\| \int_0^t A_1 M_{\eta,\eta}(T-r) A_{-1} [G(z(r)) - G(w(r))] \, dr \right\|^p$$

and

$$I_2 = \mathbf{E} \left\| \int_0^t M_{\eta,\eta}(T-r) [\tilde{h}(r, z(r)) - \tilde{h}(r, w(r))] \, dW(r) \right\|^p.$$

Using Hölder’s inequality, we obtain

$$I_1 \leq \left( \int_0^t \|M_{\eta,\eta}(T-r)\|_{H^1}^{p/(p-1)} \, dr \right)^{p-1} \int_0^t \mathbf{E} \|G(z(r)) - G(w(r))\|_{H^{-1}}^p \, dr$$

$$\begin{aligned}
 &\leq \left( \int_0^t C_\eta^{p/(p-1)} (T-r)^{-\eta p/(\alpha(p-1))} dr \right)^{p-1} \\
 &\quad \times \int_0^t C_2^p \left( \max_{t \in [0, T]} \mathbf{E} \|z(t)\|^p + \max_{t \in [0, T]} \mathbf{E} \|w(t)\|^p \right) \mathbf{E} \|z(r) - w(r)\|^p dr \\
 &\leq C_\eta^p C_2^p \left( \max_{t \in [0, T]} \mathbf{E} \|z(t)\|^p + \max_{t \in [0, T]} \mathbf{E} \|w(t)\|^p \right) \\
 &\quad \times \left( \int_0^t (T-r)^{-\eta p/(\alpha(p-1))} dr \right)^{p-1} \int_0^t \mathbf{E} \|z(r) - w(r)\|^p dr \\
 &\leq C_\eta^p C_2^p \left( \max_{t \in [0, T]} \mathbf{E} \|z(t)\|^p + \max_{t \in [0, T]} \mathbf{E} \|w(t)\|^p \right) \\
 &\quad \times T^{p(1-\eta/\alpha)-1} \left[ \frac{p-1}{p(1-\frac{\eta}{\alpha})-1} \right]^{p-1} \int_0^t \mathbf{E} \|z(r) - w(r)\|^p dr. \tag{9}
 \end{aligned}$$

Using the Burkholder–Davis–Gundy inequality from Lemma 2, we find that

$$\begin{aligned}
 I_2 &\leq \kappa(p) \mathbf{E} \left[ \left( \int_0^t \|M_{\eta, \eta}(T-r)[\hbar(r, z(r)) - \hbar(r, w(r))]\|_{L_0^2}^2 dr \right)^{p/2} \right] \\
 &\leq \kappa(p) \left( \int_0^t \|M_{\eta, \eta}(T-r)\|^{2p/(p-2)} dr \right)^{(p-2)/2} \\
 &\quad \times \int_0^t \mathbf{E} \|\hbar(r, z(r)) - \hbar(r, w(r))\|_{L_0^2}^p dr \\
 &\leq \kappa(p) C_\eta^p L_2^p \int_0^t \mathbf{E} \|z(r) - w(r)\|^p dr. \tag{10}
 \end{aligned}$$

Using (9) and (10) in (8), we infer that

$$\mathbf{E} \|v^\lambda(t, z) - v^\lambda(t, w)\|^p \leq \frac{C_v}{\lambda^p} \int_0^t \mathbf{E} \|z(r) - w(r)\|^p dr,$$

where

$$\begin{aligned}
 C_v &= 2^{p-1} \|C^*\|^p C_\eta^{2p} \left\{ C_2^p \left( \max_{t \in [0, T]} \mathbf{E} \|z(t)\|^p + \max_{t \in [0, T]} \mathbf{E} \|w(t)\|^p \right) T^{p[1-\eta/\alpha]-1} \right. \\
 &\quad \left. \times \left[ \frac{p-1}{p[1-\frac{\eta}{\alpha}]-1} \right]^{p-1} + \kappa(p) L_2^p \right\}.
 \end{aligned}$$

Since inequality (7) can be obtained in a similar manner, we omit its proof here. □

For any  $\lambda > 0$ , we define the operator  $\mathcal{F}_\lambda(z(t)) : L^p(\Sigma, H^\beta) \rightarrow L^p(\Sigma, H^\beta)$  by

$$\begin{aligned} \mathcal{F}_\lambda(z(t)) := & M_\eta(t)z_0 + \int_0^t (t-r)^{\eta-1}M_{\eta,\eta}(t-r)[G(z(r)) + Cv^\lambda(r, z)] dr \\ & + \int_0^t (t-r)^{\eta-1}M_{\eta,\eta}(t-r)\tilde{h}(r, z(r)) dW(r). \end{aligned}$$

**Theorem 1.** *If conditions (i) and (ii) are satisfied, then the time-fractional SNSE (5) has a unique mild solution  $(z(t))_{t \in [0, T]}$  in  $L^p(\Sigma, H^\beta)$  for  $p \geq 2$ ,  $\eta p \neq 1$ , and  $0 \leq \beta < \alpha < 2$ .*

To prove this result, we use the Banach contraction principle to demonstrate that the operator  $\mathcal{F}_\lambda(z(t))$  has a fixed point, which is a mild solution to (5). To this end, we first prove the following lemma.

**Lemma 7.** *For  $p \geq 2$ ,  $0 \leq \beta < \alpha < 2$ , and for any  $z \in L^p(\Sigma, H^\beta)$ , the operator  $\mathcal{F}_\lambda(z(t))$  is continuous on  $[0, T]$  in the  $L^p$  sense.*

*Proof.* For  $0 \leq \tau_1 < \tau_2 \leq T$  and a fixed  $z \in L^p(\Sigma, H^\beta)$ , we have

$$\begin{aligned} & \mathbf{E} \left\| \mathcal{F}_\lambda(z(\tau_2)) - \mathcal{F}_\lambda(z(\tau_1)) \right\|_{H^\beta}^p \\ & \leq 4^{p-1} \left[ \mathbf{E} \left\| (M_\eta(\tau_2) - M_\eta(\tau_1))z_0 \right\|_{H^\beta}^p \right. \\ & \quad + \mathbf{E} \left\| \int_0^{\tau_2} (\tau_2-r)^{\eta-1}M_{\eta,\eta}(\tau_2-r)G(z(r)) dr - \int_0^{\tau_1} (\tau_1-r)^{\eta-1}M_{\eta,\eta}(\tau_1-r)G(z(r)) dr \right\|_{H^\beta}^p \\ & \quad + \mathbf{E} \left\| \int_0^{\tau_2} (\tau_2-r)^{\eta-1}M_{\eta,\eta}(\tau_2-r)Cv^\lambda(r, z) dr - \int_0^{\tau_1} (\tau_1-r)^{\eta-1}M_{\eta,\eta}(\tau_1-r)Cv^\lambda(r, z) dr \right\|_{H^\beta}^p \\ & \quad + \mathbf{E} \left\| \int_0^{\tau_2} (\tau_2-r)^{\eta-1}M_{\eta,\eta}(\tau_2-r)\tilde{h}(r, z(r)) dW(r) \right. \\ & \quad \left. - \int_0^{\tau_1} (\tau_1-r)^{\eta-1}M_{\eta,\eta}(\tau_1-r)\tilde{h}(r, z(r)) dW(r) \right\|_{H^\beta}^p \left. \right] \\ & \leq 4^{p-1} \sum_{j=1}^4 J_j, \end{aligned} \tag{11}$$

where

$$J_1 = \mathbf{E} \left\| (M_\eta(\tau_2) - M_\eta(\tau_1))z_0 \right\|_{H^\beta}^p,$$

$$\begin{aligned}
 J_2 &= \mathbf{E} \left\| \int_0^{\tau_2} (\tau_2 - r)^{\eta-1} M_{\eta,\eta}(\tau_2 - r) G(z(r)) \, dr \right. \\
 &\quad \left. - \int_0^{\tau_1} (\tau_1 - r)^{\eta-1} M_{\eta,\eta}(\tau_1 - r) G(z(r)) \, dr \right\|_{H^\beta}^p \\
 &\leq 3^{p-1} \left[ \mathbf{E} \left\| \int_0^{\tau_1} (\tau_1 - r)^{\eta-1} (M_{\eta,\eta}(\tau_2 - r) - M_{\eta,\eta}(\tau_1 - r)) G(z(r)) \, dr \right\|_{H^\beta}^p \right. \\
 &\quad \left. + \mathbf{E} \left\| \int_0^{\tau_1} [(\tau_2 - r)^{\eta-1} - (\tau_1 - r)^{\eta-1}] M_{\eta,\eta}(\tau_2 - r) G(z(r)) \, dr \right\|_{H^\beta}^p \right. \\
 &\quad \left. + \mathbf{E} \left\| \int_{\tau_1}^{\tau_2} (\tau_2 - r)^{\eta-1} M_{\eta,\eta}(\tau_2 - r) G(z(r)) \, dr \right\|_{H^\beta}^p \right] \\
 &= 3^{p-1} (J_{21} + J_{22} + J_{23}),
 \end{aligned}$$

$$\begin{aligned}
 J_3 &= \mathbf{E} \left\| \int_0^{\tau_2} (\tau_2 - r)^{\eta-1} M_{\eta,\eta}(\tau_2 - r) C v^\lambda(r, z) \, dr \right. \\
 &\quad \left. - \int_0^{\tau_1} (\tau_1 - r)^{\eta-1} M_{\eta,\eta}(\tau_1 - r) C v^\lambda(r, z) \, dr \right\|_{H^\beta}^p \\
 &\leq 3^{p-1} \left[ \mathbf{E} \left\| \int_0^{\tau_1} (\tau_1 - r)^{\eta-1} (M_{\eta,\eta}(\tau_2 - r) - M_{\eta,\eta}(\tau_1 - r)) C v^\lambda(r, z) \, dr \right\|_{H^\beta}^p \right. \\
 &\quad \left. + \mathbf{E} \left\| \int_0^{\tau_1} [(\tau_2 - r)^{\eta-1} - (\tau_1 - r)^{\eta-1}] M_{\eta,\eta}(\tau_2 - r) C v^\lambda(r, z) \, dr \right\|_{H^\beta}^p \right. \\
 &\quad \left. + \mathbf{E} \left\| \int_{\tau_1}^{\tau_2} (\tau_2 - r)^{\eta-1} M_{\eta,\eta}(\tau_2 - r) C v^\lambda(r, z) \, dr \right\|_{H^\beta}^p \right] \\
 &= 3^{p-1} (J_{31} + J_{32} + J_{33}),
 \end{aligned}$$

and

$$\begin{aligned}
 J_4 &= \mathbf{E} \left\| \int_0^{\tau_2} (\tau_2 - r)^{\eta-1} M_{\eta,\eta}(\tau_2 - r) \tilde{h}(r, z(r)) \, dW(r) \right. \\
 &\quad \left. - \int_0^{\tau_1} (\tau_1 - r)^{\eta-1} M_{\eta,\eta}(\tau_1 - r) \tilde{h}(r, z(r)) \, dW(r) \right\|_{H^\beta}^p
 \end{aligned}$$

$$\begin{aligned} &\leq 3^{p-1} \left[ \mathbf{E} \left\| \int_0^{\tau_1} (\tau_1 - r)^{\eta-1} (M_{\eta,\eta}(\tau_2 - r) - M_{\eta,\eta}(\tau_1 - r)) \mathfrak{h}(r, z(r)) \, dW(r) \right\|_{H^\beta}^p \right. \\ &\quad + \mathbf{E} \left\| \int_0^{\tau_1} [(\tau_2 - r)^{\eta-1} - (\tau_1 - r)^{\eta-1}] M_{\eta,\eta}(\tau_2 - r) \mathfrak{h}(r, z(r)) \, dW(r) \right\|_{H^\beta}^p \\ &\quad \left. + \mathbf{E} \left\| \int_{\tau_1}^{\tau_2} (\tau_2 - r)^{\eta-1} M_{\eta,\eta}(\tau_2 - r) \mathfrak{h}(r, z(r)) \, dW(r) \right\|_{H^\beta}^p \right] \\ &= 3^{p-1} (J_{41} + J_{42} + J_{43}). \end{aligned}$$

Using Lemma 4, we see that

$$J_1 \leq C_{\alpha\beta}^p (\tau_2 - \tau_1)^{p\eta\beta/\alpha} \mathbf{E} \|z_0\|^p.$$

Next, using Lemmas 3 and 4, condition (i), and Hölder’s inequality, we find that

$$\begin{aligned} J_{21} &= \mathbf{E} \left\| \int_0^{\tau_1} (\tau_1 - r)^{\eta-1} A_\beta [M_{\eta,\eta}(\tau_2 - r) - M_{\eta,\eta}(\tau_1 - r)] G(z(r)) \, dr \right\|^p \\ &\leq C_{\eta\beta}^p (\tau_2 - \tau_1)^{p\eta(\beta+1)\alpha} \left( \int_0^{\tau_1} (\tau_1 - r)^{(\eta-1)p/(p-1)} \, dr \right)^{p-1} \\ &\quad \times \int_0^{\tau_1} \mathbf{E} \|G(z(r))\|_{H^{-1}}^p \, dr \\ &\leq C_{\eta\beta}^p C_1^p T^{\eta p} \left[ \frac{p-1}{\eta p - 1} \right]^{p-1} \sup_{t \in [0, T]} \mathbf{E} \|z(t)\|^{2p} (\tau_2 - \tau_1)^{p\eta(\beta+1)\alpha}, \\ J_{22} &= \mathbf{E} \left\| \int_0^{\tau_1} [(\tau_2 - r)^{\eta-1} - (\tau_1 - r)^{\eta-1}] A_\beta M_{\eta,\eta}(\tau_2 - r) G(z(r)) \, dr \right\|^p \\ &\leq C_\eta^p \left( \int_0^{\tau_1} \{ [(\tau_2 - r)^{\eta-1} - (\tau_1 - r)^{\eta-1}] (\tau_2 - r)^{(\beta+1)\eta/\alpha} \}^{p/(p-1)} \, dr \right)^{p-1} \\ &\quad \times \int_0^{\tau_1} \mathbf{E} \|G(z(r))\|_{H^{-1}}^p \, dr \\ &\leq 2C_\eta^p C_1^p T \left\{ \frac{p-1}{p(\eta - \frac{\eta(\beta+1)}{\alpha}) - 1} \right\}^{p-1} \sup_{t \in [0, T]} \mathbf{E} \|z(t)\|^{2p} (\tau_2 - \tau_1)^{p\eta(\alpha-\beta-1)-\alpha/\alpha}, \end{aligned}$$

and

$$\begin{aligned}
 J_{23} &= \mathbf{E} \left\| \int_{\tau_1}^{\tau_2} (\tau_2 - r)^{\eta-1} M_{\eta,\eta}(\tau_2 - r) G(z(r)) \, dr \right\|_{H^\beta}^p \\
 &\leq C_\eta^p \left( \int_{\tau_1}^{\tau_2} [(\tau_2 - r)^{\eta-1-\eta(\beta+1)/\alpha}]^{p/(p-1)} \, dr \right)^{p-1} \int_{\tau_1}^{\tau_2} \mathbf{E} \|G(z(r))\|_{H^{-1}}^p \, dr \\
 &\leq C_\eta^p C_1^p \left\{ \frac{p-1}{p[\eta - \frac{\eta(\beta+1)}{\alpha}] - 1} \right\}^{p-1} \sup_{t \in [0, T]} \mathbf{E} \|z(t)\|^{2p} (\tau_2 - \tau_1)^{p\eta(\alpha-\beta-1)/\alpha}.
 \end{aligned}$$

Next, using Lemmas 3, 4, and 6, we obtain

$$\begin{aligned}
 J_{31} &= \mathbf{E} \left\| \int_0^{\tau_1} (\tau_1 - r)^{\eta-1} [M_{\eta,\eta}(\tau_2 - r) - M_{\eta,\eta}(\tau_1 - r)] C v^\lambda(r, z) \, dr \right\|_{H^\beta}^p \\
 &\leq C_{\eta\beta}^p (\tau_2 - \tau_1)^{p\eta\beta/\alpha} \left( \int_0^{\tau_1} (\tau_1 - r)^{(\eta-1)p/(p-1)} \, dr \right)^{p-1} \int_0^{\tau_1} \mathbf{E} \|C v^\lambda(r, z)\|^p \, dr \\
 &\leq C_{\eta\beta}^p \|C\|^p \frac{C_v}{\lambda^p} T^{\eta p} \left[ \frac{p-1}{\eta p - 1} \right]^{p-1} \left[ 1 + \int_0^t \mathbf{E} \|z(r)\|^p \, dr \right] (\tau_2 - \tau_1)^{p\eta\beta/\alpha},
 \end{aligned}$$

$$\begin{aligned}
 J_{32} &= \mathbf{E} \left\| \int_0^{\tau_1} [(\tau_2 - r)^{\eta-1} - (\tau_1 - r)^{\eta-1}] M_{\eta,\eta}(\tau_2 - r) C v^\lambda(r, z) \, dr \right\|_{H^\beta}^p \\
 &\leq C_\eta^p \left( \int_0^{\tau_1} \{ [(\tau_2 - r)^{\eta-1} - (\tau_1 - r)^{\eta-1}] (\tau_2 - r)^{(\beta+1)\eta/\alpha} \}^{p/(p-1)} \, dr \right)^{p-1} \\
 &\quad \times \int_0^{\tau_1} \mathbf{E} \|C v^\lambda(r, z)\|^p \, dr \\
 &\leq C_\eta^p \|C\|^p \frac{C_v}{\lambda^p} T \left\{ \frac{p-1}{p[\eta - \frac{\eta(\beta+1)}{\alpha}] - 1} \right\}^{p-1} \left[ 1 + \int_0^t \mathbf{E} \|z(r)\|^p \, dr \right] \\
 &\quad \times (\tau_2 - \tau_1)^{p\eta(\alpha-\beta-1)-\alpha/\alpha},
 \end{aligned}$$

and

$$J_{33} = \mathbf{E} \left\| \int_{\tau_1}^{\tau_2} (\tau_2 - r)^{\eta-1} M_{\eta,\eta}(\tau_2 - r) C v^\lambda(r, z) \, dr \right\|_{H^\beta}^p$$

$$\begin{aligned} &\leq C_{\eta}^p \left( \int_{\tau_1}^{\tau_2} [(\tau_2 - r)^{\eta-1-\eta(\beta+1)/\alpha}]^{p/(p-1)} dr \right)^{p-1} \int_{\tau_1}^{\tau_2} \mathbf{E} \|Cv^{\lambda}(r, z)\|^p dr \\ &\leq C_{\eta}^p \|C\|^p \frac{C_v}{\lambda^p} \left\{ \frac{p-1}{p[\eta - \frac{\eta(\beta+1)}{\alpha}] - 1} \right\}^{p-1} \left[ 1 + \int_0^t \mathbf{E} \|z(r)\|^p dr \right] \\ &\quad \times (\tau_2 - \tau_1)^{p\eta(\alpha-\beta-1)/\alpha}. \end{aligned}$$

Using Lemma 2 and condition (ii), we get

$$\begin{aligned} J_{41} &= \mathbf{E} \left\| \int_0^{\tau_1} (\tau_1 - r)^{\eta-1} [M_{\eta,\eta}(\tau_2 - r) - M_{\eta,\eta}(\tau_1 - r)] \tilde{h}(r, z(r)) dW(r) \right\|_{H^{\beta}}^p \\ &\leq \kappa(p) \mathbf{E} \left( \int_0^{\tau_1} \|(\tau_1 - r)^{\eta-1} A_{\beta} [M_{\eta,\eta}(\tau_2 - r) - M_{\eta,\eta}(\tau_1 - r)]\|^2 \|\tilde{h}(r, z(r))\|_{L_0^2}^2 dr \right)^{p/2} \\ &\leq \kappa(p) C_{\eta\beta}^p (\tau_2 - \tau_1)^{p\eta\beta/\alpha} \left( \int_0^{\tau_1} (\tau_1 - r)^{2p(\eta-1)/(p-2)} dr \right)^{(p-2)/2} \int_0^{\tau_1} \mathbf{E} \|\tilde{h}(r, z(r))\|_{L_0^2}^p dr \\ &\leq \kappa(p) C_{\eta\beta}^p L_1^p T^{(2p\eta-p-1)/2} \left[ \frac{p-1}{2p\eta-p-2} \right]^{p-1} \left[ 1 + \sup_{t \in [0, T]} \mathbf{E} \|z(t)\|^p \right] (\tau_2 - \tau_1)^{p\eta\beta/\alpha}, \end{aligned}$$

$$\begin{aligned} J_{42} &= \mathbf{E} \left\| \int_0^{\tau_1} [(\tau_2 - r)^{\eta-1} - (\tau_1 - r)^{\eta-1}] M_{\eta,\eta}(\tau_2 - r) \tilde{h}(r, z(r)) dW(r) \right\|_{H^{\beta}}^p \\ &\leq \kappa(p) \mathbf{E} \left( \int_0^{\tau_1} \|[(\tau_2 - r)^{\eta-1} - (\tau_1 - r)^{\eta-1}] A_{\beta} M_{\eta,\eta}(\tau_2 - r)\|^2 \|\tilde{h}(r, z(r))\|_{L_0^2}^2 dr \right)^{p/2} \\ &\leq \kappa(p) C_{\eta}^p \left( \int_0^{\tau_1} \{[(\tau_2 - r)^{\eta-1} - (\tau_1 - r)^{\eta-1}] (\tau_2 - r)^{-\eta\beta/\alpha}\}^{2p/(p-2)} dr \right)^{(p-2)/2} \\ &\quad \times \int_0^{\tau_1} \mathbf{E} \|\tilde{h}(r, z(r))\|_{L_0^2}^p dr \\ &\leq \kappa(p) C_{\eta}^p L_1^p T \left[ \frac{\alpha(p-2)}{2p\eta(\alpha-\beta) - (p+2)\alpha} \right]^{(p-2)/2} \left[ 1 + \sup_{t \in [0, T]} \mathbf{E} \|z(t)\|^p \right] \\ &\quad \times (\tau_2 - \tau_1)^{(2p\eta(\alpha-\beta) - (p+2)\alpha)/(2\alpha)}, \end{aligned}$$

and

$$\begin{aligned}
 J_{43} &= \mathbf{E} \left\| \int_{\tau_1}^{\tau_2} (\tau_2 - r)^{\eta-1} M_{\eta,\eta}(\tau_2 - r) \hbar(r, z(r)) \, dW(r) \right\|_{H^\beta}^p \\
 &\leq \kappa(p) \mathbf{E} \left( \int_{\tau_1}^{\tau_2} \|(\tau_2 - r)^{\eta-1} A_\beta M_{\eta,\eta}(\tau_2 - r)\|^2 \|\hbar(r, z(r))\|_{L_0^2}^2 \, dr \right)^{p/2} \\
 &\leq \kappa(p) C_\eta^p \left( \int_{\tau_1}^{\tau_2} [(\tau_2 - r)^{\eta-1-\eta\beta/\alpha}]^{2p/(p-2)} \, dr \right)^{(p-2)/2} \int_{\tau_1}^{\tau_2} \mathbf{E} \|\hbar(r, z(r))\|_{L_0^2}^p \, dr \\
 &\leq \kappa(p) C_\eta^p L_1^p \left[ \frac{\alpha(p-2)}{2p\eta(\alpha-\beta) - (p+2)\alpha} \right]^{(p-2)/2} \left[ 1 + \sup_{t \in [0, T]} \mathbf{E} \|z(t)\|^p \right] \\
 &\quad \times (\tau_2 - \tau_1)^{(2p\eta(\alpha-\beta) - p\alpha)/(2\alpha)}.
 \end{aligned}$$

Combining the above inequalities and plugging them into (11), we see that

$$\mathbf{E} \|\mathcal{F}_\lambda(z(\tau_2)) - \mathcal{F}_\lambda(z(\tau_1))\|_{H^\beta}^p \rightarrow 0 \quad \text{as } \tau_2 - \tau_1 \rightarrow 0.$$

Hence  $\mathcal{F}_\lambda(z(t))$  is continuous on  $[0, T]$ . □

*Proof of Theorem 1.* We first show that  $\mathcal{F}_\lambda$  maps  $L^p(\Sigma, H^\beta)$  into  $L^p(\Sigma, H^\beta)$ . Indeed, for  $z \in L^p(\Sigma, H^\beta)$ , we have

$$\mathbf{E} \|\mathcal{F}_\lambda(z(t))\|_{H^\beta}^p \leq 4^{p-1} \left[ \sup_{t \in [0, T]} \mathbf{E} \|M_\eta(t)z_0\|_{H^\beta}^p + \sup_{t \in [0, T]} \sum_{i=1}^3 \mathbf{E} \|\Theta_i^z(t)\|_{H^\beta}^p \right],$$

where

$$\begin{aligned}
 &\sup_{t \in [0, T]} \mathbf{E} \|M_\eta(t)z_0\|_{H^\beta}^p \leq C_\alpha^p T^{-p\eta\beta/\alpha} \mathbf{E} \|z_0\|^p, \\
 &\sup_{t \in [0, T]} \mathbf{E} \|\Theta_1^z(t)\|_{H^\beta}^p \\
 &\leq \sup_{t \in [0, T]} \mathbf{E} \left\| \int_0^t (t-r)^{\eta-1} M_{\eta,\eta}(t-r) Gz(r) \, dr \right\|_{H^\beta}^p \\
 &\leq \sup_{t \in [0, T]} \mathbf{E} \left\| \int_0^t (t-r)^{\eta-1} A_{\beta+1} M_{\eta,\eta}(t-r) A_{-1} Gz(r) \, dr \right\|_{H^\beta}^p \\
 &\leq \sup_{t \in [0, T]} C_\eta^p \left( \int_0^t (t-r)^{p[\eta-1-\eta(\beta+1)/\alpha]/(p-1)} \, dr \right)^{p-1} \int_0^t \mathbf{E} \|A_{-1} G(z(r))\|^p \, dr \\
 &\leq C_\eta^p C_1^p \left\{ \frac{p-1}{p[\eta - \frac{\eta(\beta+1)}{\alpha}] - 1} \right\}^{p-1} T^{p[\eta-\eta(\beta+1)/\alpha]-1} \max_{t \in [0, T]} \mathbf{E} \|z(t)\|^2,
 \end{aligned}$$



$$\begin{aligned}
 & \sup_{t \in [0, T]} \mathbf{E} \|\Theta_2^z(t)\|_{H^\beta}^p \\
 & \leq \sup_{t \in [0, T]} \mathbf{E} \left\| \int_0^t (t-r)^{\eta-1} M_{\eta, \eta}(t-r) C v^\lambda(r, z(r)) \, dr \right\|_{H^\beta}^p \\
 & \leq \sup_{t \in [0, T]} \mathbf{E} \left\| \int_0^t (t-r)^{\eta-1} A_\beta M_{\eta, \eta}(t-r) C v^\lambda(r, z(r)) \, dr \right\|_{H^\beta}^p \\
 & \leq \sup_{t \in [0, T]} C_\eta^p \left( \int_0^t (t-r)^{p[\eta-1-\eta\beta/\alpha]/(p-1)} \, dr \right)^{p-1} \int_0^t \mathbf{E} \|C v^\lambda(r, z(r))\|^p \, dr \\
 & \leq C_\eta^p \|C\|^p \frac{C_v}{\lambda^p} \left\{ \frac{p-1}{p[\eta-\frac{\eta\beta}{\alpha}]-1} \right\}^{p-1} T^{p[\eta-\eta\beta/\alpha]} \left[ 1 + \max_{t \in [0, T]} \int_0^t \mathbf{E} \|z(r)\|^p \, dr \right],
 \end{aligned}$$

and

$$\begin{aligned}
 & \sup_{t \in [0, T]} \mathbf{E} \|\Theta_3^z(t)\|_{H^\beta}^p \\
 & \leq \sup_{t \in [0, T]} \mathbf{E} \left\| \int_0^t (t-r)^{\eta-1} M_{\eta, \eta}(t-r) \tilde{h}(r, z(r)) \, dW(r) \right\|_{H^\beta}^p \\
 & \leq \kappa(p) \sup_{t \in [0, T]} \mathbf{E} \left( \int_0^t \|(t-r)^{\eta-1} A_\beta M_{\eta, \eta}(t-r)\|^2 \|\tilde{h}(r, z(r))\|_{L_0^2}^2 \, dr \right)^{p/2} \\
 & \leq \kappa(p) C_\eta^p \left( \int_0^t (t-r)^{2p[\eta-1-\eta\beta/\alpha]/(p-2)} \, dr \right)^{(p-2)/2} \int_0^t \mathbf{E} \|\tilde{h}(r, z(r))\|_{L_0^2}^p \, dr \\
 & \leq \kappa(p) C_\eta^p L_1^p \left[ \frac{p-2}{p(2\beta-1-\frac{\eta\beta}{\alpha})-2} \right]^{(p-2)/2} T^{p(2\beta-1-\eta\beta/\alpha)-2/2} \left[ 1 + \sup_{t \in [0, T]} \mathbf{E} \|z(t)\|^p \right].
 \end{aligned}$$

Hence  $\mathbf{E} \|\mathcal{F}_\lambda(z(t))\|_{H^\beta}^p \leq \infty$ . Using the continuity of the operator  $\mathcal{F}_\lambda$ , we get that  $\mathcal{F}_\lambda z \in H^\beta$ . Hence for each  $\lambda > 0$ ,  $\mathcal{F}_\lambda$  indeed maps  $L^p(\Sigma, H^\beta)$  into itself, as asserted. Next, for  $z, w \in L^p(\Sigma, H^\beta)$ , we have

$$\begin{aligned}
 & \mathbf{E} \|(\mathcal{F}_\lambda z)(t) - (\mathcal{F}_\lambda w)(t)\|_{H^\beta}^p \\
 & \leq 3^{p-1} \left\{ \mathbf{E} \left\| \int_0^t (t-r)^{\eta-1} M_{\eta, \eta}(t-r) [Gz(r) - Gw(r)] \, dr \right\|_{H^\beta}^p \right. \\
 & \quad \left. + \mathbf{E} \left\| \int_0^t (t-r)^{\eta-1} M_{\eta, \eta}(t-r) [Cv^\lambda(r, z(r)) - Cv^\lambda(r, w(r))] \, dr \right\|_{H^\beta}^p \right.
 \end{aligned}$$

$$\begin{aligned}
 & + \mathbf{E} \left\| \int_0^t (t-r)^{\eta-1} M_{\eta,\eta}(t-r) [\tilde{h}(r, z(r)) - \tilde{h}(r, w(r))] dW(r) \right\|_{H^\beta}^p \Big\} \\
 & = J_1 + J_2 + J_3,
 \end{aligned}$$

where

$$\begin{aligned}
 J_1 & = \mathbf{E} \left\| \int_0^t (t-r)^{\eta-1} M_{\eta,\eta}(t-r) [Gz(r) - Gw(r)] dr \right\|_{H^\beta}^p \\
 & \leq \mathbf{E} \left\| \int_0^t (t-r)^{\eta-1} A_1 M_{\eta,\eta}(t-r) A_{\beta-1} [Gz(r) - Gw(r)] dr \right\|_{H^\beta}^p \\
 & \leq C_\eta^p \mathbf{E} \left( \int_0^t (t-r)^{(\eta-1-\eta/\alpha)p/(p-1)} dr \right)^{p-1} \int_0^t \mathbf{E} \|A_\beta [Gz(r) - Gw(r)]\|_{H^{-1}}^p dr \\
 & \leq C_\eta^p C_2^p \left( \max_{t \in [0,T]} \mathbf{E} \|z(t)\|_{H^\beta}^p + \max_{t \in [0,T]} \mathbf{E} \|w(t)\|_{H^\beta}^p \right) \left[ \frac{T^{(\eta-1-\eta/\alpha)p/(p-1)}}{(\eta-1-\frac{\eta}{\alpha})\frac{p}{p-1}} \right]^{p-1} \\
 & \quad \times \int_0^t \mathbf{E} \|z(r) - w(r)\|_{H^\beta}^p dr \\
 & \leq C_\eta^p C_2^p \left( \max_{t \in [0,T]} \mathbf{E} \|z(t)\|_{H^\beta}^p + \max_{t \in [0,T]} \mathbf{E} \|w(t)\|_{H^\beta}^p \right) \left[ \frac{p-1}{p\eta[1-\frac{1}{\alpha}]-1} \right]^{p-1} \\
 & \quad \times T^{p\eta[1-1/\alpha]-1} \int_0^t \mathbf{E} \|z(r) - w(r)\|_{H^\beta}^p dr, \\
 J_2 & = \mathbf{E} \left\| \int_0^t (t-r)^{\eta-1} M_{\eta,\eta}(t-r) [Cv^\lambda(r, z(r)) - Cv^\lambda(r, w(r))] dr \right\|_{H^\beta}^p \\
 & \leq \mathbf{E} \left\| \int_0^t (t-r)^{\eta-1} M_{\eta,\eta}(t-r) A_\beta [Cv^\lambda(r, z(r)) - Cv^\lambda(r, w(r))] dr \right\|_{H^\beta}^p \\
 & \leq C_\eta^p \|C\|^p \left( \int_0^t (t-r)^{(\eta-1)p/(p-1)} dr \right)^{p-1} \int_0^t \mathbf{E} \|v^\lambda(r, z(r)) - v^\lambda(r, w(r))\|_{H^\beta}^p \\
 & \leq C_\eta^p \|C\|^p T^{(p\eta-1)} \left[ \frac{p-1}{\eta p - 1} \right]^{p-1} \int_0^t \mathbf{E} \|v^\lambda(r, z(r)) - v^\lambda(r, w(r))\|_{H^\beta}^p \\
 & \leq C_\eta^p \|C\|^p \frac{C_v}{\lambda^p} T^{(p\eta)} \left[ \frac{p-1}{\eta p - 1} \right]^{p-1} \int_0^t \mathbf{E} \|z(r) - w(r)\|_{H^\beta}^p dr,
 \end{aligned}$$

and

$$\begin{aligned}
 J_3 &= \mathbf{E} \left\| \int_0^t (t-r)^{\eta-1} M_{\eta,\eta}(t-r) [\hbar(r, z(r)) - \hbar(r, w(r))] dW(r) \right\|_{H^\beta}^p \\
 &\leq \kappa(p) C_\eta^p \mathbf{E} \left( \int_0^t \|(t-r)^{\eta-1} M_{\eta,\eta}(t-r) A_\beta [\hbar(r, z(r)) - \hbar(r, w(r))]\|_{L_0^2}^2 dr \right)^{p/2} \\
 &\leq \kappa(p) C_\eta^p \left( \int_0^t (t-r)^{(\eta-1)2p/(p-2)} dr \right)^{(p-2)/2} \int_0^t \mathbf{E} \|A_\beta [\hbar(r, z(r)) - \hbar(r, w(r))]\|_{L_0^2}^p dr \\
 &\leq \kappa(p) C_\eta^p L_2^p T^{(2p\eta-p-2)/2} \left[ \frac{p-2}{2\eta p - p - 2} \right]^{(p-2)/2} \int_0^t \mathbf{E} \|z(r) - w(r)\|_{H^\beta}^p dr.
 \end{aligned}$$

Hence

$$\begin{aligned}
 &\mathbf{E} \|(\mathcal{F}_\lambda z)(t) - (\mathcal{F}_\lambda w)(t)\|_{H^\beta}^p \\
 &\leq 3^{p-1} \left[ C_\eta^p C_2^p \left( \max_{t \in [0, T]} \mathbf{E} \|z(t)\|_{H^\beta}^p + \max_{t \in [0, T]} \mathbf{E} \|w(t)\|_{H^\beta}^p \right) \right. \\
 &\quad \times \left. \left[ \frac{p-1}{p\eta \left[1 - \frac{1}{\alpha}\right] - 1} \right]^{p-1} T^{p\eta[1-1/\alpha]-1} + C_\eta^p \|C\|^p \frac{C_v}{\lambda^p} T^{(p\eta)} \left[ \frac{p-1}{\eta p - 1} \right]^{p-1} \right. \\
 &\quad \left. + \kappa(p) C_\eta^p L_2^p T^{(2p\eta-p-2)/2} \left[ \frac{p-2}{2\eta p - p - 2} \right]^{(p-2)/2} \int_0^t \mathbf{E} \|z(r) - w(r)\|_{H^\beta}^p dr. \right.
 \end{aligned}$$

As a result, there is a positive number  $\Phi(\lambda)$  such that

$$\mathbf{E} \|(\mathcal{F}_\lambda z)(t) - (\mathcal{F}_\lambda w)(t)\|_{H^\beta}^p \leq \Phi(\lambda) \int_0^t \mathbf{E} \|z(r) - w(r)\|_{H^\beta}^p dr.$$

Iterating, we see that, for any natural number  $n \geq 1$ ,

$$\mathbf{E} \|(\mathcal{F}_\lambda^n z)(t) - (\mathcal{F}_\lambda^n w)(t)\|_{H^\beta}^p \leq \frac{(T\Phi(\lambda))^n}{n!} \int_0^t \mathbf{E} \|z(r) - w(r)\|_{H^\beta}^p dr$$

for any  $\lambda > 0$ . When  $n$  is sufficiently large (so that  $(T\Phi(\lambda))^n/n! < 1$ ), using Banach’s fixed point theorem, we obtain that the operator  $\mathcal{F}_\lambda^n$  has a unique fixed point  $u_\lambda \in L^p(\Sigma, H^\beta)$ . Since

$$\mathcal{F}_\lambda^n (\mathcal{F}_\lambda(u_\lambda)) = \mathcal{F}_\lambda (\mathcal{F}_\lambda^n(u_\lambda)) = \mathcal{F}_\lambda(u_\lambda)$$

and  $\mathcal{F}_\lambda^n$  has a unique fixed point, it follows that  $\mathcal{F}_\lambda(u_\lambda)$  is the unique fixed point of  $\mathcal{F}_\lambda^n$ , that is,  $F_\lambda(u_\lambda) = u_\lambda$ , which is a mild solution to (5). □

**Remark 1.** Theorem 1 proves that under assumptions (i) and (ii), the time-fractional SNSE (5) has a unique mild solution  $(z(t))_{t \in [0, T]}$  of the form (6).

**Theorem 2.** *If conditions (i)–(iii) hold true and the functions  $\tilde{h}$  and  $G$  are uniformly bounded, then the time-fractional stochastic Navier–Stokes equation (5) is approximately controllable.*

*Proof.* Let  $z_\lambda$  be the fixed point of  $\mathcal{F}_\lambda$ . Using the stochastic Fubini theorem, one can show that the fixed point of  $\mathcal{F}_\lambda$  satisfies

$$\begin{aligned} z_\lambda(T) &= z_T - \lambda(\lambda I + \Upsilon_0^T)^{-1} [\mathbf{E}z_T - M_\eta(T)z_0] \\ &\quad + \int_0^T \lambda(\lambda I + \Upsilon_0^T)^{-1} (T - r)^{\eta-1} M_{\eta, \eta}(T - r) G(z_\lambda(r)) \, dr \\ &\quad + \int_0^T \lambda(\lambda I + \Upsilon_0^T)^{-1} [(T - r)^{\eta-1} M_{\eta, \eta}(T - r) \tilde{h}(r, z_\lambda(r)) - \varphi(r)] \, dW(r). \end{aligned}$$

Since the functions  $\tilde{h}$  and  $G$  are uniformly bounded, there exist constants  $D_1 > 0$  and  $D_2 > 0$  so that

$$\|\tilde{h}(r, z_\lambda(r))\|^p \leq D_1, \quad \text{and} \quad \|G(z_\lambda(r))\|^p \leq D_2.$$

Therefore we can find a subsequence  $\{\tilde{h}(c, z_\lambda(c)), G(z_\lambda(c))\}$ , which converges weakly to  $\{\tilde{h}(r, z(r)), G(z(r))\}$ .

Using the Lebesgue dominated convergence theorem, we obtain

$$\mathbf{E} \int_0^T \|M_{\eta, \eta}(T - r) [\tilde{h}(r, z_\lambda(r)) - \tilde{h}(r, z(r))]\|^p \, dr \rightarrow 0$$

and

$$\mathbf{E} \int_0^T \|M_{\eta, \eta}(T - r) [G(z_\lambda(r)) - G(z(r))]\|^p \, dr \rightarrow 0.$$

Hence

$$\begin{aligned} &\mathbf{E} \|z_\lambda(T) - z_T\|^p \\ &\leq 6^{p-1} \left[ \|\lambda(\lambda I + \Upsilon_0^T)^{-1} [\mathbf{E}z_T - M_\eta(T)z_0]\|^p \right. \\ &\quad + \mathbf{E} \left( \int_0^T (T - r)^{\eta-1} \|\lambda(\lambda I + \Upsilon_0^T)^{-1} \varphi(r)\|_{L_0^2}^p \, dr \right) \\ &\quad \left. + \mathbf{E} \left( \int_0^T (T - r)^{\eta-1} \|\lambda(\lambda I + \Upsilon_0^T)^{-1}\| \|M_{\eta, \eta}(T - r) [G(z_\lambda(r)) - G(z(r))]\| \, dr \right)^p \right] \end{aligned}$$

$$\begin{aligned}
 &+ \mathbf{E} \left( \int_0^T (T-r)^{\eta-1} \|\lambda(\lambda I + \mathcal{Y}_0^T)^{-1}\| \|M_{\eta,\eta}(T-r)G(z(r))\| \, dr \right)^p \\
 &+ \mathbf{E} \left( \int_0^T \|(T-r)^{\eta-1}\lambda(\lambda I + \mathcal{Y}_0^T)^{-1}M_{\eta,\eta}(T-r)[\tilde{h}(r, z_\lambda(r)) - \tilde{h}(r, z(r))]\|_{L_0^2}^2 \, dr \right)^{p/2} \\
 &+ \mathbf{E} \left( \int_0^T \|(T-r)^{\eta-1}\lambda(\lambda I + \mathcal{Y}_0^T)^{-1}M_{\eta,\eta}(T-r)\tilde{h}(r, z(r))\|_{L_0^2}^2 \, dr \right)^{p/2} \Big].
 \end{aligned}$$

From assumption (iii) it follows that the operator

$$\lambda(\lambda I + \mathcal{Y}_0^T)^{-1} \rightarrow 0 \quad \text{strongly as } \lambda \rightarrow 0.$$

In addition,

$$\|\lambda(\lambda I + \mathcal{Y}_0^T)^{-1}\| \leq 1.$$

Thus, using the Lebesgue dominated convergence theorem and the compactness of the operator  $M_{\eta,\eta}$ , we obtain

$$\mathbf{E}\|z_\lambda(T) - z_T\|^p \rightarrow 0 \quad \text{as } \lambda \rightarrow 0.$$

Hence using the definition of approximate controllability, the time-fractional SNSE (5) is approximately controllable on  $[0, T]$ , as asserted. □

**Remark 2.** Theorem 2 proves that under assumptions (i)–(iii), there exists a control function that steers the solution of the time-fractional stochastic Navier–Stokes equation (5) from the initial state  $z_0$  to the neighborhood of the final state  $z_T$  in  $[0, T]$ .

### 4 Example

To demonstrate the applicability of the theoretical results on approximate controllability, we consider the following two-dimensional time-fractional stochastic Navier–Stokes equation in the square domain  $\mathcal{O} = (0, 1)^2$  with Dirichlet boundary conditions:

$$\begin{aligned}
 &\partial_t^\eta z(t, x, y) + \nu(-\Delta)^{\alpha/2} z(t, x, y) \\
 &= Cv(t, x, y) + \sigma z(t, x, y) \frac{dW(t)}{dt} \quad \forall t \in (0, 1] \tag{12} \\
 &z(0, x, y) = z_0(x, y) = \sin(\pi x) \sin(\pi y),
 \end{aligned}$$

where  $z(t, x, y)$  denotes the unknown variable,  $\eta = 0.7$  is the fractional time derivative order,  $\alpha = 1.8$  is the fractional Laplacian order,  $\nu = 0.1$  is the viscosity coefficient,  $\sigma = 0.05$  is the noise intensity, the control operator  $C = I$ , and  $W(t)$  is a standard Wiener process. The control input is

$$v(t, x, y) = \sin(\pi x) \sin(\pi y), \quad t \in [0, 1].$$

The target state is the velocity field pattern induced by the control input, i.e., the first Fourier mode  $\sin(\pi x) \sin(\pi y)$ .

Setting  $A = \nu(-\Delta)^{\alpha/2}$ , the mild solution of (12) is given by

$$z(t) = E_{\eta}(-At^{\eta})z_0 + \int_0^t (t-s)^{\eta-1} E_{\eta,\eta}(-A(t-s)^{\eta})v(s) ds + \sigma \int_0^t (t-s)^{\eta-1} E_{\eta,\eta}(-A(t-s)^{\eta})z(s) dW(s).$$

Here  $E_{\eta,\beta}$  denotes the Mittag-Leffler function, and the last term is an Itô stochastic convolution. Under the assumptions stated in Section 2, analyticity of  $A$ , admissibility of  $C$ , linear growth, and Lipschitz condition on the noise coefficient, this integral equation admits a unique mild solution in  $L^2(\Omega; C([0, T]; H))$ .

### 4.1 Numerical scheme

1. *Fractional derivative.* We approximated the fractional derivative using the Grünwald–Letnikov formula with recursive computation of weights:

$$w_0 = 1, \quad w_k = \left(1 - \frac{\eta + 1}{k}\right)w_{k-1}, \quad k \geq 1.$$

2. *Fractional Laplacian.* We approximated the fractional Laplacian spectrally. For a velocity field  $z(x, y)$  expanded in sine basis, the fractional Laplacian is applied in Fourier space:

$$(-\Delta)^{\alpha/2}z(x, y) = \sum_{m,n} (\pi^2(m^2 + n^2))^{\alpha/2} \hat{z}_{m,n} \sin(m\pi x) \sin(n\pi y),$$

where  $\hat{z}_{m,n}$  are the Fourier coefficients.

3. *Noise term.* Discretized as

$$\sigma z^n \frac{\Delta W^n}{\Delta t}, \quad \Delta W^n \sim \mathcal{N}(0, \Delta t),$$


with independent realizations at each spatial point and time step.

4. *Time integration.* Semi-implicit Euler scheme is used:

$$z^{n+1} = z^n + \Delta t(-\nu(-\Delta)^{\alpha/2}z^n + v^n + \text{noise} - \text{fractional derivative term}).$$

### 4.2 Results and interpretation

Figure 2 shows the evolution of the  $L^2$  norm of the velocity field  $\|z(t, \cdot, \cdot)\|_{L^2}$  with and without control over time  $t \in [0, 1]$ . This clearly demonstrates that the controlled trajectory of the velocity field approaches the desired target pattern  $\sin(\pi x) \sin(\pi y)$  over time. Despite the presence of stochastic perturbations and memory effects, the applied



**Figure 2.** Time evolution of the  $L^2$  norm of the velocity field  $\|z(t, \cdot, \cdot)\|_{L^2}$  under the controlled and uncontrolled stochastic fractional Navier–Stokes dynamics.

control successfully guides the system close to the intended state. This illustrates the approximate controllability of the time-fractional stochastic Navier–Stokes system, confirming the theoretical results derived in Section 3.

**Author contributions.** All authors (R.C., S.R., and J.J.N.) have contributed as follows: methodology, R.C.; formal analysis, R.C.; validation, R.C., S.R., and J.J.N.; writing – original draft preparation, R.C.; writing – review & editing, S.R. and J.J.N. All authors have read and approved the published version of the manuscript.

**Conflicts of interest.** The authors declare no conflicts of interest.

## References

1. S. Barnett, *Introduction to Mathematical Control Theory*, Oxford Appl. Math. Comput. Sci. Ser., Clarendon Press, Oxford, 1975.
2. A. Bensoussan, R. Temam, Équations stochastiques du type Navier–Stokes, *J. Funct. Anal.*, **13**(2):195–222, 1973, [https://doi.org/10.1016/0022-1236\(73\)90045-1](https://doi.org/10.1016/0022-1236(73)90045-1).
3. Z. Brzeźniak, E. Hausenblas, J. Zhu, 2D stochastic Navier–Stokes equations driven by jump noise, *Nonlinear Anal., Theory Methods Appl.*, **79**:122–139, 2013, <https://doi.org/10.1016/j.na.2012.10.011>.
4. M. Caputo, Vibrations of an infinite viscoelastic layer with a dissipative memory, *J. Acoust. Soc. Am.*, **56**(3):897–904, 1974, <https://doi.org/10.1121/1.1903344>.
5. R.F. Curtain, H. Zwart, *An Introduction to Infinite-Dimensional Linear Systems Theory*, Texts Appl. Math., Vol. 21, Springer, New York, 1995, <https://doi.org/10.1007/978-1-4612-4224-6>.

6. G. Da Prato, A. Debussche, Two-dimensional Navier–Stokes equations driven by a space–time white noise, *J. Funct. Anal.*, **196**(1):180–210, 2002, <https://doi.org/10.1006/jfan.2002.3919>.
7. J.P. Dauer, N.I. Mahmudov, Controllability of stochastic semilinear functional differential equations in Hilbert spaces, *J. Math. Anal. Appl.*, **290**(2):373–394, 2004, <https://doi.org/10.1016/j.jmaa.2003.09.069>.
8. P.M. de Carvalho-Neto, G. Planas, Mild solutions to the time fractional Navier–Stokes equations in  $\mathbb{R}^N$ , *J. Differ. Equations*, **259**(7):2948–2980, 2015, <https://doi.org/10.1016/j.jde.2015.04.008>.
9. L. Debbi, Well-posedness of the multidimensional fractional stochastic Navier–Stokes equations on the torus and on bounded domains, *J. Math. Fluid Mech.*, **18**(1):25–69, 2016, <https://doi.org/10.1007/s00021-015-0234-5>.
10. K. Diethelm, *The Analysis of Fractional Differential Equations: An Application-Oriented Exposition Using Differential Operators of Caputo Type*, Lect. Notes Math., Vol. 2004, Springer, Berlin, 2010, <https://doi.org/10.1007/978-3-642-14574-2>.
11. X.-L. Ding, J.J. Nieto, X. Wang, Analytical solutions for fractional partial delay differential-algebraic equations with Dirichlet boundary conditions defined on a finite domain, *Fract. Calc. Appl. Anal.*, **25**(2):408–438, 2022, <https://doi.org/10.1007/s13540-022-00021-7>.
12. B.-S. Han,  $L_p$ -solvability and Hölder regularity for stochastic time fractional Burgers’ equations driven by multiplicative space-time white noise, *Stoch. Partial Differ. Equ. Anal. Comput.*, **13**(1):26–79, 2025, <https://doi.org/10.1007/s40072-024-00329-w>.
13. M. Hofmanová, R. Zhu, X. Zhu, Global existence and non-uniqueness for 3D Navier–Stokes equations with space-time white noise, *Arch. Ration. Mech. Anal.*, **247**(3):46, 2023.
14. R.E. Kalman, Mathematical description of linear dynamical systems, *J. Soc. Ind. Appl. Math., Ser. A, Control*, 1:152–192, 1963.
15. K. Karthikeyan, D. Tamizharasan, J.J. Nieto, K.S. Nisar, Controllability of second-order differential equations with state-dependent delay, *IMA J. Math. Control Inform.*, **38**(4):1072–1083, 2021, <https://doi.org/10.1093/imamci/dnab027>.
16. A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North-Holland Math. Stud., Vol. 204, Elsevier, Amsterdam, 2006.
17. J.U. Kim, Strong solutions of the stochastic Navier-Stokes equations in  $\mathbb{R}^3$ , *Indiana Univ. Math. J.*, **59**(5):1853–1886, 2010, <https://doi.org/10.1512/iumj.2010.59.3930>.
18. Raphael Kruse, *Strong and Weak Approximation of Semilinear Stochastic Evolution Equations*, Lect. Notes Math., Vol. 2093, Springer, Cham, 2014, <https://doi.org/10.1007/978-3-319-02231-4>.
19. M. Kwaśnicki, Ten equivalent definitions of the fractional Laplace operator, *Fract. Calc. Appl. Anal.*, **20**(1):7–51, 2017, <https://doi.org/10.1515/fca-2017-0002>.
20. L.D. Landau, E.M. Lifshitz, *Fluid Mechanics*, Course of Theoretical Physics, Vol. 6, Pergamon Press, Oxford, 1959. Translated from the Russian by J.B. Sykes and W.H. Reid.
21. J. Liao, F. Sueur, P. Zhang, Global controllability of the Navier–Stokes equations in the presence of curved boundary with no-slip conditions, *J. Math. Fluid Mech.*, **24**(3):71, 2022, <https://doi.org/10.1007/s00021-022-00689-0>.



22. J.T. Machado, V. Kiryakova, F. Mainardi, Recent history of fractional calculus, *Commun. Nonlinear Sci. Numer. Simul.*, **16**(3):1140–1153, 2011, <https://doi.org/10.1016/j.cnsns.2010.05.027>.
23. S. Momani, Z. Odibat, Analytical solution of a time-fractional Navier–Stokes equation by Adomian decomposition method, *Appl. Math. Comput.*, **177**(2):488–494, 2006, <https://doi.org/10.1016/j.amc.2005.11.025>.
24. I. Podlubny, *Fractional Differential Equations: An Introduction to Fractional Derivatives, Fractional Differential Equations, to Methods of Their Solution and Some of Their Applications*, Math. Sci. Eng., Vol. 198, Academic Press, San Diego, CA, 1999.
25. R. Sakthivel, S. Suganya, S.M. Anthoni, Approximate controllability of fractional stochastic evolution equations, *Comput. Math. Appl.*, **63**(3):660–668, 2012, <https://doi.org/10.1016/j.camwa.2011.11.024>.
26. V. Singh, R. Chaudhary, L. Kumar Som, Approximate controllability of stochastic differential system with non-Lipschitz conditions, *Stochastic Anal. Appl.*, **40**(3):505–519, 2022, <https://doi.org/10.1080/07362994.2021.1930050>.
27. X.-X. Xi, M. Hou, X.-F. Zhou, Y. Wen, Approximate controllability for mild solution of time-fractional Navier–Stokes equations with delay, *Z. Angew. Math. Phys.*, **72**(3):113, 2021, <https://doi.org/10.1007/s00033-021-01542-6>.
28. L. Xu, T. Shen, X. Yang, J. Liang, Analysis of time fractional and space nonlocal stochastic incompressible Navier–Stokes equation driven by white noise, *Comput. Math. Appl.*, **78**(5):1669–1680, 2019, <https://doi.org/10.1016/j.camwa.2018.12.022>.
29. Y. Zhou, L. Peng, On the time-fractional Navier–Stokes equations, *Comput. Math. Appl.*, **73**(6):874–891, 2017, <https://doi.org/10.1016/j.camwa.2016.03.026>.
30. G.-A. Zou, B. Wang, Stochastic Burgers’ equation with fractional derivative driven by multiplicative noise, *Comput. Math. Appl.*, **74**(12):3195–3208, 2017, <https://doi.org/10.1016/j.camwa.2017.08.023>.
31. G.A. Zou, G. Lv, J.-L. Wu, Stochastic Navier–Stokes equations with Caputo derivative driven by fractional noises, *J. Math. Anal. Appl.*, **461**(1):595–609, 2018, <https://doi.org/10.1016/j.jmaa.2018.01.027>.